

FORM-PTO-1390
Rev. 12-29-99)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371**

001560-403

U.S. APPLICATION NO. (If known, see 37 C.F.R. 1.5)

To be assigned

09/869360

INTERNATIONAL APPLICATION NO.
PCT/JP00/07706

INTERNATIONAL FILING DATE
01 November 2000

PRIORITY DATE CLAIMED
01 November 1999

TITLE OF INVENTION

BLOOD VESSEL LIPID DEPOSITION-PREVENTIVE AGENT COMPRISING CHYMASE-INHIBITOR

APPLICANT(S) FOR DO/EO/US

Harukazu FUKAMI and Hidenori URATA

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and the PCT Articles 22 and 39(1).
4. ☐ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
- ☒ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
- ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
- ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
- ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
- ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern other document(s) or information included:

11. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
 ☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:

International Search Report

Copy of cover page from the published PCT application (WO 01/32214)

Copy of Japanese Request Form

PCT Notice Informing the Applicant of the Communication of the International Application to the Designated Offices (Form PCT/IB/308)

U.S. APPLICATION NO. [redacted] (see 37 CFR 1.50)

INTERNATIONAL APPLICATION NO.

ATTORNEY'S DOCKET NUMBER

To be assigned

09/869360

PCT/JP00/07706

001560-403

17. ☒ The following fees are submitted:

CALCULATIONS

PTO USE ONLY

Basic National Fee (37 CFR 1.492(a)(1)-(5)):

Neither international preliminary examination fee (37 CFR 1.482)
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO
and International Search Report not prepared by the EPO or JPO \$1,000.00 (960)

International preliminary examination fee (37 CFR 1.482) not paid to
USPTO but International Search Report prepared by the EPO or JPO \$860.00 (970)

International preliminary examination fee (37 CFR 1.482) not paid to USPTO
but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$710.00 (958)

International preliminary examination fee paid to USPTO (37 CFR 1.482)
but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00 (956)

International preliminary examination fee paid to USPTO (37 CFR 1.482)
and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00 (962)

ENTER APPROPRIATE BASIC FEE AMOUNT =

\$ 860.00

Surcharge of \$130.00 (154) for furnishing the oath or declaration later than
months from the earliest claimed priority date (37 CFR 1.492(e)).

20 ☐ 30 ☐

\$

Claims	Number Filed	Number Extra	Rate
Total Claims	10 -20 =	0	X\$18.00 (966)

\$

Independent Claims	3 -3 =	0	X\$80.00 (964)
--------------------	--------	---	----------------

\$

Multiple dependent claim(s) (if applicable)			+ \$270.00 (968)
---	--	--	------------------

\$

TOTAL OF ABOVE CALCULATIONS =

\$ 860.00

Reduction for 1/2 for filing by small entity, if applicable (see below).

\$

SUBTOTAL =

\$ 860.00

Processing fee of \$130.00 (156) for furnishing the English translation later than
months from the earliest claimed priority date (37 CFR 1.492(f)).

20 ☐ 30 ☐

\$

+

TOTAL NATIONAL FEE =

\$ 860.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by
an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 (581) per property +

\$ 40.00

TOTAL FEES ENCLOSED =

\$ 900.00

Amount to be:
refunded

\$

charged

\$

- a. ☐ Small entity status is hereby claimed.
- b. ☒ A check in the amount of \$ 900.00 to cover the above fees is enclosed.
- c. ☐ Please charge my Deposit Account No. 02-4800 in the amount of \$ to cover the above fees. A duplicate copy of this sheet is enclosed.
- d. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 02-4800. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Ronald L. Grudziecki
BURNS, DOANE, SWECKER & MATHIS, L.L.P.
P.O. Box 1404
Alexandria, Virginia 22313-1404
(703) 836-6620

SIGNATURE

Donna M. Meuth

NAME

Date: June 28, 200136,607

REGISTRATION NUMBER

09/869360

JC18 Rec'd PCT/PTO 2 8 JUN 2001

Patent

Attorney's Docket No. 001560-403

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of)
)
Harukazu FUKAMI et al) Group Art Unit: To be assigned
)
Application No.: U.S. National Phase of) Examiner: To be assigned
PCT/JP00/07706, filed November 1, 2000)
)
Filed: June 28, 2001)
)
For: BLOOD VESSEL LIPID)
DEPOSITION-PREVENTIVE)
AGENT COMPRISING CHYMASE-)
INHIBITOR)

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231


Sir:

Prior to the examination of the above-identified patent application, please enter the following amendments.

IN THE CLAIMS

Kindly replace claims 6-7 as follows:

6. (Amended) The preventive or therapeutic agent according to claim 1,
wherein said chymase inhibitor is a quinazoline derivative represented by the formula (1):



wherein, the ring A represents an aryl ring,

R¹ represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower alkyl group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;

R² and R³, which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 4 carbons that may be substituted, a halogen atom, a hydroxy group, a


lower alkoxy group having 1 to 4 carbons, an amino group, a lower alkylamino group having 1 to 4 carbons that may be substituted, a lower aralkylamino group having 7 to 10 carbons that may be substituted, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, or a carboxylic group; or

when the ring A is a benzene ring, R¹ and R², together with the benzene ring to be substituted, may form a fused heterocyclic ring that may be substituted with a carboxylic acid, and a carbon atom in said fused heterocyclic ring may form a carbonyl group wherein R³ is as defined above; and

X represents a hydrogen atom, a lower alkyl group having 1 to 4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;

or a pharmaceutically acceptable salt thereof.

7. (Amended) The pharmaceutical composition according to claim 3, wherein said chymase inhibitor is a quinazoline derivative represented by the formula (1):



wherein, the ring A represents an aryl ring,

R¹ represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower alkyl group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;


R² and R³, which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 4 carbons that may be substituted, a halogen atom, a hydroxy group, a lower alkoxy group having 1 to 4 carbons, an amino group, a lower alkylamino group having 1 to 4 carbons that may be substituted, a lower aralkylamino group having 7 to 10 carbons that may be substituted, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, or a carboxylic group; or

when the ring A is a benzene ring, R¹ and R², together with the benzene ring to be substituted, may form a fused heterocyclic ring that may be substituted with a carboxylic acid, and a carbon atom in said fused heterocyclic ring may form a carbonyl group wherein R³ is as defined above; and

X represents a hydrogen atom, a lower alkyl group having 1-4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;
or a pharmaceutically acceptable salt thereof.

Please add new claims 9-10 as follows.

--9. The preventive or therapeutic agent according to claim 2, wherein said chymase inhibitor is a quinazoline derivative represented by the formula (1):



wherein, the ring A represents an aryl ring,

R¹ represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower alkyl group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;


R² and R³, which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 4 carbons that may be substituted, a halogen atom, a hydroxy group, a lower alkoxy group having 1 to 4 carbons, an amino group, a lower alkylamino group having 1 to 4 carbons that may be substituted, a lower aralkylamino group having 7 to 10 carbons that may be substituted, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, or a carboxylic group; or

when the ring A is a benzene ring, R¹ and R², together with the benzene ring to be substituted, may form a fused heterocyclic ring that may be substituted with a carboxylic acid, and a carbon atom in said fused heterocyclic ring may form a carbonyl group wherein R³ is as defined above; and

X represents a hydrogen atom, a lower alkyl group having 1 to 4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;

or a pharmaceutically acceptable salt thereof.

10. The pharmaceutical composition according to claim 4, wherein said chymase inhibitor is a quinazoline derivative represented by the formula (1):



wherein, the ring A represents an aryl ring,

R¹ represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower alkyl

group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;

R^2 and R^3 , which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 4 carbons that may be substituted, a halogen atom, a hydroxy group, a lower alkoxy group having 1 to 4 carbons, an amino group, a lower alkylamino group having 1 to 4 carbons that may be substituted, a lower aralkylamino group having 7 to 10 carbons that may be substituted, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, or a carboxylic group; or

when the ring A is a benzene ring, R^1 and R^2 , together with the benzene ring to be substituted, may form a fused heterocyclic ring that may be substituted with a carboxylic acid, and a carbon atom in said fused heterocyclic ring may form a carbonyl group wherein R^3 is as defined above; and

X represents a hydrogen atom, a lower alkyl group having 1-4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;

or a pharmaceutically acceptable salt thereof.--

REMARKS


Prior to examination of the above-identified application, entry of the foregoing, and consideration of the above amendments are respectfully requested.

Claims 6-7 have been amended simply to delete multiple dependencies. New claims 9-10 have been added, in view of deletion of the multiple dependencies. It is respectfully submitted that no new matter has been added by the above amendments.

In the event that there are any questions relating to this Preliminary Amendment, or to the application in general, it would be appreciated if the Examiner would telephone the undersigned attorney at 508-339-3684 concerning such questions so that prosecution of this application may be expedited.

Respectfully submitted,

BURNS, DOANE, SWECKER & MATHIS, L.L.P.

By: 
Donna M. Meuth
Registration No. 36,607


P.O. Box 1404
Alexandria, Virginia 22313-1404
(703) 836-6620

Date: June 28, 2001

Attachment to Preliminary Amendment dated June 28, 2001

Marked-up Claims 6-7

6. (Amended) The preventive or therapeutic agent according to claim 1 [or 2], wherein said chymase inhibitor is a quinazoline derivative represented by the formula (1):



wherein, the ring A represents an aryl ring,

R¹ represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower alkyl

Attachment to Preliminary Amendment dated June 28, 2001

Marked-up Claims 6-7

group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;

R^2 and R^3 , which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 4 carbons that may be substituted, a halogen atom, a hydroxy group, a lower alkoxy group having 1 to 4 carbons, an amino group, a lower alkylamino group having 1 to 4 carbons that may be substituted, a lower aralkylamino group having 7 to 10 carbons that may be substituted, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, or a carboxylic group; or


when the ring A is a benzene ring, R^1 and R^2 , together with the benzene ring to be substituted, may form a fused heterocyclic ring that may be substituted with a carboxylic acid, and a carbon atom in said fused heterocyclic ring may form a carbonyl group wherein R^3 is as defined above; and

Attachment to Preliminary Amendment dated June 28, 2001

Marked-up Claims 6-7

X represents a hydrogen atom, a lower alkyl group having 1 to 4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;
or a pharmaceutically acceptable salt thereof.

7. (Amended) The pharmaceutical composition according to claim 3 [or 4], wherein said chymase inhibitor is a quinazoline derivative represented by the formula (1):



wherein, the ring A represents an aryl ring,

R¹ represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic

Marked-up Claims 6-7

R^2 and R^3 , which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 4 carbons that may be substituted, a halogen atom, a hydroxy group, a lower alkoxy group having 1 to 4 carbons, an amino group, a lower alkylamino group having 1 to 4 carbons that may be substituted, a lower aralkylamino group having 7 to 10 carbons that may be substituted, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino

Attachment to Preliminary Amendment dated June 28, 2001

Marked-up Claims 6-7

group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, or a carboxylic group; or

when the ring A is a benzene ring, R¹ and R², together with the benzene ring to be substituted, may form a fused heterocyclic ring that may be substituted with a carboxylic acid, and a carbon atom in said fused heterocyclic ring may form a carbonyl group wherein R³ is as defined above; and

X represents a hydrogen atom, a lower alkyl group having 1-4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;

or a pharmaceutically acceptable salt thereof.

001560-403

3/PRTS

09/869360
JC18 Rec'd PCT/PTO 2 8 JUN 2001
STY-H848

- 1 -

DESCRIPTION

BLOOD VESSEL LIPID DEPOSITION-PREVENTIVE
AGENT COMPRISING CHYMASE-INHIBITOR

5

TECHNICAL FIELD

The present invention provides a preventive or
therapeutic agent for diseases accompanied by abnormal
vascular function in which lipid deposition in the blood
vessel is involved, a preventive or therapeutic
pharmaceutical composition for diseases accompanied by
abnormal vascular function, and a suppressing agent of
lipid deposition in the blood vessel.

BACKGROUND ART

A major mechanism of lipid deposition in the blood
vessel is believed that monocytes and macrophages
infiltrate into the injured vascular endothelial cells,
and thereby these cells incorporate oxygenated low
density lipoproteins (LDL) in excess and turn into the
so-called foam cells that have accumulated droplets of
cholesterol esters (Ross R., Nature 362: 801, 1993). It
is thought that foam cells, together with T cells and
vascular smooth muscle cells, form fatty streaks, and the
interaction between the cells facilitates pathological
processes, generating vascular lesions such as
arteriosclerosis including atherosclerosis.

In many epidemiological studies in recent years,
hyperlipemia has been defined as a risk factor of
arteriosclerosis, and in fact various drugs that regulate
blood levels of lipids such as cholesterol and
triglyceride have been reported. For example, drugs such
as Plavastatin that suppress cholesterol biosynthesis by
inhibiting 3-hydroxy-3-methylglutaryl coenzyme A (HMG-
CoA) have been widely used. These drugs can indeed lower
lipid levels in the blood during the administration
period, but various problems have been pointed out: once
the administration is suspended the level returns to the

level before administration; the effect is not adequate in patients with severe high-cholesterolemia; or improvement in blood lipid levels does not always lead to life lengthening.

5 These drugs are also known to be associated with side effects such as myopathy and abnormal hepatic function, and are likely to inhibit the biosynthesis of physiological components such as ubiquinone and dolichol, raising a possibility to elicit an adverse effect. Other
10 therapeutic agents of hyperlipemia include drugs that influence lipoprotein metabolism in the blood vessel such as Clofibrate, drugs that suppress the absorption of cholesterol from the intestinal tract such as Nicomol and Colestyramine, and the like. None of them, however, are
15 satisfactory in terms of efficacy and side effects, and thus there is a need for the development of further excellent drugs in terms of efficacy and safety.

 On the other hand, chymase is a serine protease that is widely distributed in the tissue such as the skin, the
20 heart, vascular walls, intestinal tracts, etc. as a granular component in mast cell (Mast Cell Proteases in Immunology and Biology; Caughey, G.H., Ed; Marcel Dekker, Inc.: New York, 1995). Chymase is known to participate in a synthetic process independent of angiotensin
25 converting enzyme in the conversion of angiotensin I to angiotensin II.

 It is also reported that in the aorta of atherosclerosis or arterial aneurysm a chymase-dependent angiotensin II (AngII) forming activity was observed to
30 be significantly higher than in the aorta without atherosclerosis or arterial aneurysm (M. Ihara, et al., Hypertension 32: 514-20, 1998) and that the expression of chymase mRNA is increased in the aorta of monkeys that were fed a high-cholesterol diet for 6 months (S. Takai, et al., FEBS Lett. 412: 86-90, 1997).
35

 It has also been indicated that LDL can be restrictively degraded by chymase, and that the

modified LDL binds to mast cell granules (Mast Cell
Proteases in Immunology and Biology; Caughey, G.H., Ed;
Marcel Dekker, Inc.: New York, 1995). LDL-granule
complex is likely to be easily incorporated into
5 macrophages. These clinical findings and experimental
results implicate the involvement of intravascular
chymase in atheroma formation, but the role of chymase in
pathological and physiological states has not been
elucidated and the study to clarify this point has just
10 begun. In recent years, search for substances that
inhibit chymase activity are underway in addition to the
elucidation of physiological actions of chymase.

As chymase inhibitors, there have been reported: a
low molecular weight chymase inhibitor described in a
15 textbook (Protease Inhibitors; Barrett et al., Eds.:
Elsevier Science B.V.: Amsterdam, 1996); reported as a
peptidyl inhibitor, α -keto acid derivative (WO 93-25574,
Proc. Natl. Acad. Sci. USA 92: 6738, 1995), α,α -
difluoro- β -keto acid derivative (Japanese Unexamined
20 Patent Publication (Kokai) No. 9-124691), a tripeptide
inhibitor (WO 93-03625), and a phosphoric acid derivative
(Oleksyszyn et al., Biochemistry 30: 485, 1991); as
peptide-like inhibitors, a trifluoromethylketone
derivative (WO 96-33974, Japanese Unexamined Patent
25 Publication (Kokai) No. 10-53579) and an acetamide
derivative (Japanese Unexamined Patent Publication
(Kokai) No. 10-7661, Japanese Unexamined Patent
Publication (Kokai) No. 10-53579, Japanese Unexamined
Patent Publication (Kokai) No. 11-246437, WO 99-41277, WO
30 98-18794, WO 96-39373); as non-peptidyl inhibitors, a
triazine derivative (Japanese Unexamined Patent
Publication (Kokai) No. 8-208654, Japanese Unexamined
Patent Publication (Kokai) No. 10-245384), a phenolester
derivative (Japanese Unexamined Patent Publication
35 (Kokai) No. 10-87567), a cephem derivative (Japanese
Unexamined Patent Publication (Kokai) No. 10-87493), an

isoxazole derivative (Japanese Unexamined Patent Publication (Kokai) No. 11-1479), an imidazolidine derivative (WO 96-04248), a hydantoin derivative (Japanese Unexamined Patent Publication (Kokai) No. 9-31061), a quinazoline derivative (WO 97-11941), and the like. However, no drugs or therapeutic regimens have been established that employ the inhibition of chymase activity as a therapeutic strategy.

DISCLOSURE OF THE INVENTION

The present invention intends to provide a safe preventive or therapeutic agent that suppresses the progression of pathological processes in abnormal vascular function in which lipid deposition is involved and prevents the progression of various complications, which will improve the quality of life of the patients.

After intensive study to resolve the above problems, the inventors of the present invention have created an animal model of arterial lipid deposition induced by a high-cholesterol diet, obtained a finding that, surprisingly, a chymase inhibitor suppresses lipid deposition in the blood vessel and improves abnormal vascular function, and clarified the association of chymase activity with lipid deposition, and thereby have completed the present invention.

Thus, the present invention relates to a preventive or therapeutic agent for diseases accompanied by abnormal vascular function in which lipid deposition in the blood vessel is involved, said agent comprising a chymase inhibitor as an active ingredient.

The present invention also relates to a preventive or therapeutic pharmaceutical composition for diseases accompanied by abnormal vascular function, wherein the chymase inhibitor is blended at an amount that suppresses lipid deposition in the blood vessel.

Furthermore, the present invention relates to a suppressing agent of lipid deposition in the blood vessel, comprising a chymase inhibitor as an active

ingredient.

BRIEF EXPLANATION OF DRAWINGS

Figure 1 is a graph showing that arterial lipid deposition was increased in the hamsters that received a high-cholesterol diet (HC) as compared to the hamsters that received a normal diet (NC), and that the lipid deposition was decreased by the administration of chymase inhibitor (HC + compound 18 of the present invention).

Figure 2 is a graph showing the correlation between the plasma levels of total cholesterol (A) or LDL-cholesterol (B) and the level of chymase-like activity in a high-cholesterol diet model.

Figure 3 is a graph showing chymase activity in transgenic (Tg) mice in which human chymase was excessively expressed.

Figure 4 is a graph showing that arterial lipid deposition in the human chymase Tg mice was significantly higher than that of the control mice when a high cholesterol diet was given.


EMBODIMENT FOR CARRYING OUT THE INVENTION

As used herein, diseases accompanied by abnormal vascular function in which lipid deposition in the blood vessel is involved include diseases that result from abnormal vascular function caused by lipid deposition in the blood vessel, diseases of which symptoms are aggravated by the development of abnormal vascular function caused by lipid deposition in the blood vessel, diseases of which cure is delayed by the development of abnormal vascular function caused by lipid deposition in the blood vessel and the like. For example, the onset of abnormal vascular function associated with lipid deposition in the blood vessel is observed in arteriosclerosis, cardiac acute coronary syndrome such as unstable angina, and acute myocardial infarction, restenosis after percutaneous transluminal coronary angioplasty, obstructive arteriosclerosis, obstructive thrombotic vasculitis, atherosclerosis, cerebral

infarction, intermittent claudication, lower limb gangrene, renal vascular hypertension, renal arterial aneurysm, renal infarction, and the like.

Chymase inhibitors suitable for use in the present invention can be selected as substances that can inhibit chymase activity by using methods capable of being performed by a person skilled in the art. As a method of selection, there can be mentioned a method described in Example 1 below. The compounds obtained in this way include, for example, a low molecular weight chymase inhibitor described in a textbook (Protease Inhibitors; Barrett et al., Ed.: Elsevier Science B.V.: Amsterdam, 1996); reported as a peptidyl inhibitor, α -keto acid derivative (WO 93-25574, Proc. Natl. Acad. Sci. USA 92: 6738, 1995), α,α -difluoro- β -keto acid derivative (Japanese Unexamined Patent Publication (Kokai) No. 9-124691), a tripeptide inhibitor (WO 93-03625), and a phosphoric acid derivative (Oleksyszyn et al., Biochemistry 30: 485, 1991); as a peptide-like inhibitor, a trifluoromethylketone derivative (WO 96-33974, Japanese Unexamined Patent Publication (Kokai) No. 10-53579) and an acetamide derivative (Japanese Unexamined Patent Publication (Kokai) No. 10-7661, Japanese Unexamined Patent Publication (Kokai) No. 10-53579, Japanese Unexamined Patent Publication (Kokai) No. 11-246437, WO 99-41277, WO 98-18794, WO 96-39373); as non-peptidyl inhibitors, a triazine derivative (Japanese Unexamined Patent Publication (Kokai) No. 8-208654, Japanese Unexamined Patent Publication (Kokai) No. 10-245384), a phenolester derivative (Japanese Unexamined Patent Publication (Kokai) No. 10-87567), a cephem derivative (Japanese Unexamined Patent Publication (Kokai) No. 10-87493), an isoxazole derivative (Japanese Unexamined Patent Publication (Kokai) No. 11-1479), an imidazolidine derivative (WO 96-04248), a hydantoin derivative (Japanese Unexamined Patent Publication (Kokai) No. 9-

31061), a quinazoline derivative (WO 97-11941), and the like, and as a representative example of preferred chymase inhibitors, there can be mentioned a compound represented by the following formula (1):



10 wherein, the ring A represents an aryl ring,

R^1 represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower alkyl group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;

R^2 and R^3 , which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 4 carbons that may be substituted, a halogen atom, a hydroxy group, a lower alkoxy group having 1 to 4 carbons, an amino group, a lower alkylamino group having 1 to 4 carbons that may be substituted, a lower

aralkylamino group having 7 to 10 carbons that may be substituted, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, or a carboxylic group; or

when the ring A is a benzene ring, R^1 and R^2 , together with the benzene ring to be substituted, may form a fused heterocyclic ring that may be substituted with a carboxylic acid, and a carbon atom in said fused heterocyclic ring may form a carbonyl group wherein R^3 is as defined above; and

X represents a hydrogen atom, a lower alkyl group having 1 to 4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;

and a pharmaceutically acceptable salt thereof.

As a preferred example of the aryl ring represented by the ring A in the general formula (1), a benzene ring or a naphthalene ring is illustrated.

As a preferred example of R^1 which is a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group or a lower aralkylamino group having 7 to 12 carbons that may be substituted with a carboxylic group, there can be illustrated a methylamino group, an ethylamino group, a propylamino group, a butylamino group, a carboxymethylamino group, a carboxyethylamino group, a

carboxypropylamino group, a carboxybutylamino group, a
benzylamino group, a phenethylamino group, a
phenylpropylamino group, a phenylbutylamino group, a
carboxybenzylamino group, a carboxyphenethylamino group,
5 a carboxyphenylpropylamino group, a
carboxyphenylbutylamino group, and the like.

As a preferred example of R^1 which is an amino group
acylated with a lower fatty acid having 1 to 4 carbons
that may be substituted with a carboxylic group, an amino
10 group acylated with an aromatic carboxylic acid that may
be substituted with a carboxylic group, or an amino group
acylated with a heteroaromatic carboxylic acid that may
be substituted with a carboxylic group, there can be
illustrated a formylamino group, an acetylamino group, a
15 propionylamino group, a butyrylamino group, a
benzoylamino group, a naphthoylamino group, a
pyridinecarbonylamino group, a pyrrolecarbonylamino
group, a carboxyacetylamino group, a
carboxypropionylamino group, a carboxybutyrylamino group,
20 a carboxylbenzoylamino group, a carboxynaphthoylamino
group, a carboxypyridinecarbonylamino group, a
carboxypyrrolecarbonylamino group, and the like.

As a preferred example of R^1 which is an amino group
sulfonylated with a lower alkanesulfonic acid having 1 to
25 4 carbons that may be substituted with a carboxylic
group, an amino group sulfonylated with an aromatic
sulfonic acid that may be substituted with a carboxylic
group, or an amino group sulfonylated with a
heteroaromatic sulfonic acid that may be substituted with
30 a carboxylic group, there can be illustrated a
methanesulfonylamino group, an ethanesulfonylamino group,
a propanesulfonylamino group, a butanesulfonylamino
group, a benzenesulfonylamino group, a
naphthalenesulfonylamino group, a pyridinesulfonylamino
35 group, a pyrrolesulfonylamino group, a
carboxymethanesulfonylamino group, a
carboxyethanesulfonylamino group, a

carboxypropanesulfonylamino group, a
carboxybutanesulfonylamino group, a
carboxybenzenesulfonylamino group, a
carboxynaphthalenesulfonylamino group, a
5 carboxypyridinesulfonylamino group, a
carboxypyrrolesulfonylamino group, and the like.

As a preferred example of R^1 which is a lower alkyl
group having 1 to 4 carbons substituted with a carboxylic
group, there can be illustrated an acetic group,
10 propionic group, butyric group, valeric group, and the
like. As a preferred example of R^1 which is a lower
alkylene group having 2 to 4 carbons substituted with a
carboxylic group, there can be illustrated an acrylic
group, a crotonic group, and the like.

15 As a preferred example of R^2 or R^3 which is a lower
alkyl group having 1 to 4 carbons that may be
substituted, there can be illustrated a straight-chain
alkyl group such as a methyl group, an ethyl group, a n-
propyl group, and a n-butyl group, a branched-chain alkyl
20 group such as an isopropyl group, a sec-butyl group, and
a t-butyl group, and as a preferred example of the
substituent of a lower alkyl group having 1 to 4 carbons,
there can be illustrated a carboxylic group, a halogen
atom such as fluorine and chlorine, a lower alkoxy group
25 having 1 to 4 carbons, an amino group, a methylamino
group, a dimethylamino group, a carboxymethylamino group,
and a carboxyethylamino group and the like. As a
preferred example of R^2 or R^3 which is a halogen atom,
there can be illustrated fluorine, chlorine, and iodine.

30 As a preferred example of R^2 or R^3 which is a lower
alkoxy group having 1 to 4 carbons, there can be
illustrated a straight-chain alkyloxy group such as a
methoxy group, an ethoxy group, a n-propyloxy group and a
n-butoxy group, a branched-chain alkyloxy group such as
35 an isopropyloxy group, a sec-butoxy group and a t-butoxy
group.

As a preferred example of R^2 or R^3 which is a lower

alkylamino group having 1 to 4 carbons that may be substituted, there can be illustrated a methylamino group, an ethylamino group, a propylamino group, a butylamino group, and the like, and as a preferred example of the substituent of the lower alkylamino group having 1 to 4 carbons, there can be illustrated a carboxylic group, a halogen atom such as fluorine and chlorine, a lower alkoxy group having 1 to 4 carbons, and the like.

As a preferred example of R^2 or R^3 which is a lower aralkylamino group having 7 to 12 carbons that may be substituted, there can be illustrated a benzylamino group, a phenethylamino group, a phenylpropylamino group, a phenylbutylamino group, and the like, and as a preferred example of the substituent of the aralkylamino group, there can be illustrated a carboxylic group, a halogen atom such as fluorine and chlorine, a lower alkoxy group having 1 to 4 carbons, and the like.

As a preferred example of R^2 or R^3 which is an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, or an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, there can be illustrated a formylamino group, an acetylamino group, a propionylamino group, a butyrylamino group, a benzoylamino group, a naphthoylamino group, a pyridinecarbonylamino group, a pyrrolocarbonylamino group, a carboxyacetylaminogroup, a carboxypropionylamino group, a carboxybutyrylamino group, a carboxylbenzoylamino group, a carboxynaphthoylamino group, a carboxypyridinecarbonylamino group, a carboxypyrrolocarbonylamino group, and the like.

As a preferred example of R^2 or R^3 which is an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a

carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, or an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, there can be illustrated a methanesulfonylamino group, an ethanesulfonylamino group, a propanesulfonylamino group, a benzenesulfonylamino group, a naphthalenesulfonylamino group, a pyridinesulfonylamino group, a pyrrolesulfonylamino group, a carboxymethanesulfonylamino group, a carboxyethanesulfonylamino group, a carboxypropanesulfonylamino group, a carboxybenzenesulfonylamino group, a carboxynaphthalenesulfonylamino group, a carboxypyridinesulfonylamino group, a carboxypyrrolesulfonylamino group, and the like.

When the ring A is a benzene ring, as a preferred example of a fused heterocyclic ring that is formed by R^1 and R^2 together with the benzene ring to be substituted, that may be substituted with a carboxylic acid, and whose carbon atom in the fused heterocyclic ring may form a carbonyl group, there can be mentioned a tetrahydroquinoline ring and a benzoxadine ring, and specifically there can be illustrated tetrahydroquinoline, benzoxadine, quinoxaline, benzodioxane, carboxytetrahydroquinoline, carboxybenzoxadine, carboxyquinoxaline, carboxybenzodioxane, and the like.

As a preferred example of X which is a lower alkyl group having 1 to 4 carbons, there can be illustrated a straight-chain alkyl group such as a methyl group, an ethyl group, a n-propyl group and a n-butyl group, and a branched-chain alkyl group such as an isopropyl group, a sec-butyl group and a t-butyl group. As a preferred example of X which is a lower alkoxy group having 1 to 4 carbons, there can be illustrated a straight-chain alkyloxy group such as a methoxy group, an ethoxy group,


a n-propoxy group and a n-butoxy group, and a branched-chain alkyloxy group such as an isopropoxyloxy group, a sec-butoxy group and a t-butoxy group. As a preferred example of X which is a halogen atom, there can be illustrated fluorine, chlorine, bromine, or iodine.

As an example of pharmaceutically acceptable salts, there can be illustrated an acid addition salt such as a hydrochlorate, a methanesulfonate, a trifluoroacetate and a nitrate, and an alkali metal salt such as a sodium salt and a potassium salt.

The quinazoline derivative represented by the formula (1) of the present invention may be synthesized according to, for example, the synthetic method (A) or (B) shown below.

Synthetic method (A)


A compound represented by the formula (2):



wherein, the ring A is as defined above, R^1 represents R^1 that may be protected with a protecting group, R^2 represents R^2 that may be protected with a protecting group, R^3 represents R^3 that may be protected with a protecting group, and R^1 , R^2 and R^3 are as defined above, is reacted with an anthranilic acid derivative represented by the formula (3):



wherein, X' represents X that may be protected with a protecting group, and X is as defined above, using a method, for example, described in Japanese Unexamined Patent Publication (Kokai) No. 6-199839, so as to obtain a sulfonylurea derivative represented by the formula (4):



wherein, A, R^{1'}, R^{2'}, R^{3'}, and X' are as defined above, and then using a condensation agent such as 1,1'-carbonyldiimidazole (hereinafter referred to as CDI), the quinazoline ring is closed, and if desired the protecting group of R¹, R², R³ or X is deprotected. In this reaction, when R¹, R² or R³ represents a group that contains a hydroxyl group, an amino group or a carboxylic group, R¹, R² or R³ may be protected, if necessary, with a protecting group such as a benzyloxycarbonyl group, a t-butoxycarbonyl group, a benzyl group, an allyl group and a t-butyl group. When X represents a hydroxyl group or an amino group, it may be protected, if necessary, with a protecting group such as a benzyloxycarbonyl group, a t-butoxycarbonyl group, a benzyl group, an allyl group or a t-butyl group.

As a compound represented by the formula (2) for use in the present invention, a commercially available or known compound or a compound that can be synthesized by a known method may be used. For example, by the synthetic method described in EP 0269141, those that can be synthesized from the corresponding sulfonamide derivative using chlorosulfonyl isocyanate can be used. For example, 3-

allyloxycarbonylmethylbenzenesulfonylisocyanate, 4-allyloxycarbonylmethylbenzenesulfonylisocyanate, 4-allyloxybenzenesulfonylisocyanate, and the like can be used.

As an anthranilic acid derivative represented by the formula (3) for use in the present invention, a commercially available or known compound or a compound that can be synthesized by a known method may be used.

For example, anthranilic acid, 4-chloroanthranilic acid, 4-methoxyanthranilic acid, 5-chloroanthranilic acid, 4-hydroxyanthranilic acid, and the like can be used.


The reaction of closing a quinazoline ring from a sulfonylurea derivative represented by the formula (4) may be performed in an aprotic solvent, for example, an ethereal solvent such as tetrahydrofuran and dioxane, a halogenic solvent such as methylene chloride, or dimethylformamide, at a temperature between -50 and 50°C, preferably at a temperature between -20°C and room temperature. For the cyclization reaction, a conventional dehydrocondensation agent such as CDI, dicyclohexylcarbodiimide (DCC), and a related carbodiimide compound, and a mixed acid anhydride can also be used. For the deprotection reaction, hydrolysis with an acid or an alkali, reduction, oxidation, etc. may be selected as appropriate.

Synthetic method (B)

A compound represented by the formula (5):




wherein, ring A, R^{1'}, R^{2'} and R^{3'} are as defined above, is condensed with an anthranilic acid derivative represented by the formula (6):



wherein, X' is as defined above, Ph represents a phenyl group, R⁴ represents a protecting group of a carboxyl group, specifically a group that can be eliminated by hydrolysis or hydrogenolysis and that can form an ester group in combination with a carboxyl group,

for example a methyl group, an ethyl group or a benzyl group, using, for example, 1,8-diazabicyclo[5,4,0]-7-undecene (hereinafter referred to as DBU), so as to obtain a compound represented by the formula (7):



wherein, ring A, R^{1'}, R^{2'}, R^{3'}, R⁴ and X' are as defined above, which is then converted by alkali hydrolysis or hydrogenolysis into the corresponding carboxylic acid represented by the formula (4), and then, as in the synthetic method (A), the quinazoline ring is closed and if desired the protecting group of R¹, R², R³ and X is deprotected. In this reaction, when R¹, R² or R³ represents a group that contains a hydroxyl group, an amino group or a carboxylic group, R¹, R² or R³ may be protected, if necessary, with a protecting group such as a benzyloxycarbonyl group, a t-butoxycarbonyl group, a benzyl group, an allyl group or a t-butyl group. When X represents a hydroxyl group or an amino group, it may be protected, if necessary, with a protecting group such as a benzyloxycarbonyl group, a t-butoxycarbonyl group, a benzyl group, an allyl group or a t-butyl group.

As a compound represented by the formula (5) for use in the present invention, a commercially available or known compound or a compound that can be synthesized by a known method may be used. For example, there can be used 3-hydroxybenzenesulfonamide, 2-aminobenzenesulfonamide, 3-aminobenzenesulfonamide, 4-aminobenzenesulfonamide, (±)-2-(4-aminosulfonylphenyl)butyrate, 3-benzyloxycarbonylamino-4-chlorobenzenesulfonamide, 4-benzyloxycarbonylamino-3-chlorobenzenesulfonamide, 4-amino-3,5-dichlorobenzenesulfonamide, 3-benzyloxycarbonylamino-4-methylbenzenesulfonamide, 4-t-

butoxycarbonyl-3-hydroxybenzenesulfonamide, 3-
benzyloxycarbonylamino-4-t-
butoxycarbonylbenzenesulfonamide, 4-t-butoxycarbonyl-3-
hydroxybenzenesulfonamide, 3-t-butoxycarbonyl-4-
5 hydroxybenzenesulfonamide, 3-acetamide-4-
methoxybenzenesulfonamide, 3-(3-
aminosulfonyl)phenylacrylic acid t-butylester, 3-amino-4-
methoxybenzenesulfonamide, 4-methoxy-3-
methylsulfonylaminobenzenesulfonamide, 3-carboxy-4-
10 hydroxy-2-naphthalenesulfonamide, 4-
benzyloxycarbonylamino-3-t-
butoxycarbonylbenzenesulfonamide, (\pm)-3-t-butoxycarbonyl-
2-oxo-1H,3H-quinoline-7-sulfonamide, (\pm)-2-t-
butoxycarbonyl-3-oxo-1,4-benzooxadine-6-sulfonamide, and
15 the like.

As an anthranilic acid derivative represented by the
formula (6) for use in the present invention, a
commercially available or known compound or a compound
that can be synthesized by a known method may be used.
20 For example, there can be used methyl 4-chloro-2-N-
phenoxycarbonylanthranilate, ethyl 4-chloro-2-N-
phenoxycarbonylanthranilate, benzyl 4-chloro-2-N-
phenoxycarbonylanthranilate, methyl 5-chloro-2-N-
phenoxycarbonylanthranilate, ethyl 5-chloro-2-N-
25 phenoxycarbonylanthranilate, benzyl 5-chloro-2-N-
phenoxycarbonylanthranilate, methyl 4-methoxy-2-N-
phenoxycarbonylanthranilate, ethyl 4-methoxy-2-N-
phenoxycarbonylanthranilate, benzyl 4-methoxy-2-N-
phenoxycarbonylanthranilate, methyl 4-hydroxy-2-N-
30 phenoxycarbonylanthranilate, ethyl 4-hydroxy-2-N-
phenoxycarbonylanthranilate, benzyl 4-hydroxy-2-N-
phenoxycarbonylanthranilate, and the like.

The reaction for obtaining a sulfonylurea derivative
represented by the formula (7) by condensing a compound
35 represented by the formula (5) and an anthranilic acid
derivative represented by the formula (6) may be
performed in an aprotic solvent, for example, an ethereal

solvent such as tetrahydrofuran and dioxane, a halogenic solvent such as methylene chloride, or dimethylformamide at a temperature between -50 and 50°C, preferably at a temperature between -20°C and room temperature. As a

5 base for use in the condensation reaction, there can be used an organic strong base such as DBU, an inorganic base such as potassium carbonate, sodium carbonate, potassium hydroxide or sodium hydroxide, or a metal base such as sodium hydride.

10 In the reaction of subjecting a sulfonylurea derivative represented by the formula (7) to alkali hydrolysis or hydrogenolysis to obtain a sulfonylurea derivative represented by the formula (4), a normal hydrolysis condition or a hydrogenolysis condition for

15 esters may be used.

The above reaction may be performed by protecting the functional groups that do not participate in the reaction, and depending on the type of the protecting group, a conventional deprotection reaction such as a

20 chemical reduction is used for deprotection. For example, when the protecting group is a t-butyl group or a t-butoxycarbonyl group, trifluoroacetic acid may be used, and when it is allyl, a palladium catalyst such as tetrakis(triphenylphosphine)palladium (0) may be used.

25 A compound of the formula (1) in which R¹ is an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic

30 group, or an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, can be obtained by acylating a compound of formula (1) in which R¹ represents an amino group, using a carboxylic acid, a carboxylic chloride or a carboxylic

35 anhydride by a conventional method.

A compound of the formula (1) in which R¹ represents an amino group sulfonylated with a lower alkane sulfonic

acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, or an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, can be obtained by sulfonylating a compound of formula (1) in which R^1 represents an amino group, using a sulfonic acid, a sulfonic chloride or a sulfonic anhydride by a conventional method.

The compound obtained in the above processes may be purified by a conventional purification method such as recrystallization and column chromatography.

Also, as needed, the compound of the formula (1) obtained in the above processes may be converted to a salt by reacting it with one of various acids or bases. As an acid that can be used to convert a compound of the formula (1) to a salt, there can be mentioned an inorganic acid such as hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid and phosphoric acid, and an organic acid such as methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, trifluoroacetic acid, citric acid, lactic acid, maleic acid, fumaric acid, tartaric acid, acetic acid, adipic acid, palmitic acid and tannic acid.

As a base that can be used to convert a compound of the formula (1) to a salt, there can be mentioned sodium hydroxide, lithium hydroxide, potassium hydroxide, and the like.

Some of the compounds of formula (1) contain an asymmetric center, and from the racemates of the compound one of the optical isomers may be isolated by one or more methods. For example, there can be used:

(1) a method that employs an optically active column,

(2) a method that attains conversion to salts using an optically active acid or base and then performs recrystallization,

(3) a method that combines the above (1) and (2), and the like.

5 These compounds can be evaluated for the suppressing effect of lipid deposition on the blood vessel by the method described in the Example 4 below.

10 When a compound claimed in the present invention is used as a preventive and/or therapeutic agent for diseases in which lipid deposition in the blood vessel is involved, a preventive or therapeutic pharmaceutical composition for diseases accompanied by abnormal vascular function, or a suppressing agent of lipid deposition in the blood vessel, one or more than one compound of the present invention, for example, may be blended and formulated into a dosage form suitable for the
15 administration regimen according to a standard method. For example, for oral administration, a dosage form such as capsules, tablets, granules, fine granules, syrups, and dry syrups may be illustrated, and for parenteral administration, in addition to injections, suppositories, 20 suppositories such as vaginal suppositories, nasal drugs such as sprays, ointments, and transdermal absorptive drugs such as transdermal absorptive tapes may be illustrated.

25 The clinical dosage of the compound of the present invention varies depending on the symptom and severity of the disease, age and the presence of complications, etc., and on the pharmaceutical formulation. In the case of oral administration, 1 to 1000 mg as an active ingredient per adult, and in the case of parenteral administration, 30 one tenth to one half the dose of the oral administration may be administered. These doses can be increased or decreased, as appropriate, depending on the age of the patient and disease states, etc.

35 According to the present invention, the chymase inhibitor may be administered alone i.e. without blending with other active ingredients, but it is also possible to blend with other active ingredients and administer as

pharmaceutical compositions by taking into account indicated diseases, symptoms, complications, etc. Furthermore, combined use with other active ingredients may also be possible. The amount of the above other
5 active ingredient used is decided considering, but is not limited to, the minimum amount that exhibits effect by a single agent, development of side effects, and the like.

At the time of treatment, the selection of a pharmaceutical formulation containing the chymase
10 inhibitor alone as active ingredient and a formulation containing it with other active ingredients may be made by a physician depending on the age and symptom of the patient, etc.

The toxicity of the compound of the present
15 invention is low, and its acute toxicity value LD₅₀ against 5-week old male mice at 24 hours after oral administration is 1 g/kg or greater. The value is more than 50-times that of the expected clinical dosage, and thus it is judged that the safety of these compounds is
20 high.

EXAMPLES

The present invention will now be described more specifically based on the Examples, and it should be noted that the scope of the present invention is not
25 limited by these examples in any way.

Preparation Example 1: Synthesis of 7-chloro-3-(3-hydroxybenzenesulfonyl)-2,4(1H,3H)-quinazolinedione (compound 1):

According to the synthetic method (B), 938 mg (5.42
30 mmol) of 3-hydroxybenzenesulfonamide was dissolved in 40 ml of tetrahydrofuran, to which 892 μ l (5.96 mmol) of 1,8-diazabicyclo[5,4,0]-7-undecene (hereinafter referred to as DBU) was added dropwise. After the reaction mixture was stirred at room temperature for 15 minutes,
35 1.66 g (5.42 mmol) of methyl 4-chloro-2-N-phenoxy-carbonylanthranilate was added and stirred overnight at room temperature. An excess water was added

to the reaction mixture, which was acidified with hydrochloric acid and then extracted with ethyl acetate.

The organic layer was washed with water and saturated saline, and then dried on anhydrous magnesium sulfate and concentrated. The crude product obtained was purified by silica gel chromatography (0%-5% methanol/dichloromethane) to obtain 1.23 g (yield 59%) of methyl 4-chloro-2-[(3-hydroxybenzenesulfonylamino)carbonyl]amino}benzoate (property: colorless amorphous, PMR (δ ppm, DMSO- d_6): 3.91 (3H, s), 7.02 (1H, m), 7.09 (1H, m), 7.34 (1H, t), 7.57 (2H, m), 7.89 (1H, d), 8.38 (1H, d), 10.94 (1H, s)). Then 1.23 g (3.2 mmol) of the sulfonylurea obtained was dissolved in 20 ml of methanol, to which 10 ml of a 2N sodium hydroxide solution was added dropwise. After the reaction mixture was stirred at room temperature for 15 minutes, an excess water was added and then it was acidified with hydrochloric acid. This was stirred and the deposited crystals were taken out by filtration and dried to obtain 992 mg of a crude carboxylic acid.

The crude product obtained was dissolved in 50 ml of tetrahydrofuran (hereinafter referred to as THF), to which 434 mg (2.68 mmol) of CDI was added and stirred on ice for 30 minutes. The reaction mixture was diluted in ethyl acetate, washed with water and saturated saline, and then dried on anhydrous magnesium sulfate and concentrated to obtain a crude product. The crude product was purified by silica gel chromatography (ethyl acetate : n-hexane = 1:2) to obtain 230 mg (yield 20%: 2 steps) of the title compound. Property: colorless crystals, melting point: $>200^\circ\text{C}$ (decomposition), PMR (δ ppm, DMSO- d_6): 7.12 (2H, s), 7.24 (1H, d), 7.48 (1H, t), 7.58 (2H, s), 7.85 (1H, d), 10.28 (1H, s), 11.63 (1H, s).

Preparation Example 2: Synthesis of 3-(2-aminobenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione (compound 2):

From 2.7 g (15.7 mmol) of 2-aminobenzenesulfonamide and 4.8 g (15.7 mmol) of methyl 4-chloro-2-N-phenoxy-carbonylanthranilate in a manner similar to Preparation Example 1, 3.2 g (yield 58%: 3 steps) of the title compound was obtained. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 6.46 (2H, s), 6.65 (1H, t), 6.81 (1H, d), 7.12 (1H, s), 7.23 (1H, d), 7.34 (1H, t), 7.76 (1H, d), 7.86 (1H, d).

Preparation Example 3: Synthesis of 7-chloro-3-(2-methylsulfonylaminobenzenesulfonyl)-2,4(1H,3H)-quinazolinedione (compound 3):

Twenty-two mg (0.06 mmol) of compound 2 was dissolved in 200 μ l of pyridine, to which 11.6 μ l (0.15 mmol) of methanesulfonyl chloride was added dropwise and stirred overnight at room temperature. Excess water was added to the reaction mixture and then extracted with ethyl acetate. The organic layer was washed with an aqueous solution of 1N hydrochloric acid and saturated saline, and then dried on anhydrous magnesium sulfate and concentrated to obtain a crude product. The crude product obtained was crystallized from diethylether to obtain 16 mg (0.04 mmol) of the title compound. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 3.61 (3H, s), 7.10 (1H, d), 7.20 (1H, d), 7.74 (1H, d), 7.82-7.90 (4H, m), 8.34 (1H, d), 11.70 (1H, s).

Preparation Example 4: Synthesis of 3-(4-aminobenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione (compound 4):

From 2.7 g (15.7 mmol) of 4-aminobenzenesulfonamide and 4.8 g (15.7 mmol) of methyl 4-chloro-2-N-phenoxy-carbonylanthranilate in a manner similar to

Preparation Example 1, 7.9 g (yield 94%) of methyl 2-
{[(4-aminobenzenesulfonylamino)carbonyl]amino}-4-
chlorobenzoate was obtained. Property: colorless
amorphous, PMR (δ ppm, DMSO- d_6): 3.59 (3H, s), 5.37 (2H,
5 s), 6.45 (2H, d), 6.83 (1H, dd), 7.41 (2H, d), 7.81 (1H,
d), 8.66 (1H, d), 9.64 (1H, s).

Then from the 7.9 g (14.8 mmol) of the sulfonylurea
product, 4.3 g (yield 83%: 2 steps) of the title compound
was obtained in a similar manner. Property: colorless
10 crystals, melting point: $>200^\circ\text{C}$ (decomposition), PMR
(δ ppm, DMSO- d_6): 6.39 (2H, s), 6.63 (2H, d), 7.09 (1H,
s), 7.22 (1H, d), 7.76 (2H, d), 7.83 (1H, d), 11.51 (1H,
s).

Preparation Example 5: Synthesis of 3-(3-
15 carboxymethylbenzenesulfonyl)-7-chloro-2,4(1H,3H)-
quinazolinedione (compound 5):

After, according to the synthetic method (A), 3.27
g (11.6 mmol) of 3-allyloxycarbonylmethylbenzenesulfonyl
isocyanate was dissolved in 100 ml of anhydrous THF, 1.98
20 g (11.5 mmol) of 4-chloroanthranilic acid was added
thereto and was stirred at room temperature for 2 hours.
The reaction mixture was cooled on ice water, 1.87 g
(11.5 mmol) of CDI was added and the mixture was stirred
on ice for 30 minutes. Excess water was added to the
25 reaction mixture and then extracted with ethyl acetate.
The organic layer was washed, dried, and concentrated to
prepare a crude product, which was crystallized from a
small amount of ethyl acetate to obtain 2.0 g (yield 40%)
of 3-(3-allyloxycarbonylmethylbenzenesulfonyl)-7-chloro-
30 2,4(1H,3H)-quinazolinedione.

The above allyl product obtained was dissolved in
100 ml of a formic acid-THF (1:9) mixture, to which 700
mg of triphenylphosphine was added. The reaction vessel
was placed in the dark and the air in the reaction system
35 was displaced with nitrogen, and 700 mg of
tetrakis(triphenylphosphine)palladium (0) was added, and

stirred overnight in the dark at room temperature. The reaction mixture was concentrated under reduced pressure, and the solid obtained was washed with methylene chloride to obtain 1.47 g (yield 81%) of the title compound.

5 Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 3.76 (2H, s), 7.13 (1H, s), 7.24 (2H, d), 7.61-7.69 (2H, m), 7.86 (1H, d), 8.05 (2H, s), 12.50 (1H, br).

10 Preparation Example 6: Synthesis of 3-(4-carboxymethylbenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione (compound 6):

From 1.10 g (3.95 mmol) of 4-allyloxycarbonylmethylbenzenesulfonyl isocyanate and 678 mg (3.95 mmol) of 4-chloro-anthranilic acid in a manner similar to Preparation Example 5, 657 mg (yield 38%) of 3-(4-allyloxycarbonylbenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione was obtained. From 538 mg (1.24 mmol) of this, in a similar manner, 342 mg of the title compound (yield 70%) was obtained. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 3.75 (2H, s), 7.13 (1H, s), 7.23 (1H, d), 7.61-7.69 (2H, m), 7.86 (1H, d), 8.05 (2H, s), 12.07 (2H, br).

25 Preparation Example 7: Synthesis of (\pm)-2-{4-[(7-chloro-2,4(1H,3H)-quinazoline-3-yl)sulfonyl]phenyl}butyrate (compound 7):

From 1.02 g (3.41 mmol) of tert-butyl (\pm)-2-(4-aminosulfonylphenyl)butyrate and 1.04 g (3.41 mmol) of methyl 4-chloro-2-N-phenoxy-carbonylanthranilate in a manner similar to Preparation Example 1, 1.46 g (yield 84%) of methyl 2-[(4-[1-(tert-butoxycarbonyl)propyl]benzenesulfonylamino)carbonyl]amino]-4-chlorobenzoate (property: colorless amorphous, PMR (δ ppm, CDCl₃): 0.89 (3H, t), 1.38 (9H, s), 1.69-1.76 (1H, m), 2.03-2.10 (1H, m), 3.42 (1H, t), 3.94 (3H, s), 7.04 (1H, d), 7.47 (2H, d), 7.93 (1H, d), 8.01 (2H, d), 8.45

(1H, br), 11.04 (1H, br)) was obtained. Then, 4.3 ml (8.6 mmol) of 2N sodium hydroxide was used in a similar manner to prepare 1.43 g of a carboxylic acid. Using 463 mg (2.86 mmol) of CDI, 970 mg (yield 71%: 2 steps) of (±)-2-{4-[(7-chloro-2,4(1H,3H)-quinazoline-3-yl)sulfonyl]phenyl}butyric acid t-butylester was obtained.

The butylester product obtained was dissolved in 5 ml of dichloromethane to which 5 ml of trifluoroacetic acid was added and stirred at room temperature for 40 minutes. The reaction mixture was concentrated under reduced pressure and the crude product obtained was washed with diethylether to obtain 820 mg of the title compound (yield 96%). Property: colorless crystals, melting point: >200°C (decomposition), PMR (δppm, DMSO-d₆): 0.84 (3H, t), 1.67-1.75 (1H, m), 1.98-2.05 (1H, m), 3.62 (1H, t), 7.11 (1H, s), 7.24 (1H, d), 7.61 (2H, d), 7.86 (1H, d), 8.13 (2H, d), 11.62 (1H, s).

Preparation Example 8: Synthesis of 3-(3-amino-4-chlorobenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione (compound 8)

From 1.0 g (2.93 mmol) of 3-benzyloxycarbonylamino-4-chlorobenzenesulfonamide and 1.18 g (2.93 mmol) of 4-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example 1, 1.43 g (yield 78%) of 2-[[3-benzyloxycarbonylamino-4-chlorobenzenesulfonylamino)carbonyl]amino}-4-chlorobenzoic acid benzylester (property: colorless amorphous, PMR (δppm, DMSO-d₆): 5.19 (2H, s), 5.36 (2H, s), 7.21 (1H, dd), 7.34-7.48 (10H, m), 7.72-7.76 (2H, m), 7.97 (1H, d), 8.25 (1H, d), 8.30 (1H, d), 9.53 (1H, s), 10.30 (1H, s)) was obtained.

Among this, 1.38 g (2.20 mmol) was dissolved in 50 ml of THF, to which 200 mg of palladium-carbon (10%) was added and stirred under a hydrogen stream for 2 hours. The reaction mixture was filtered with celite to remove

palladium-carbon, and the filtrate was concentrated under reduced pressure to obtain a crude product. The crude product obtained was dissolved in 50 ml of THF, to which 356 mg (2.20 mmol) of CDI on ice was added, and, in a similar manner to Preparation Example 1, 560 mg (yield 66%: 2 steps) of the title compound was obtained.

Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 6.00 (2H, s), 7.12 (1H, s), 7.26 (2H, t), 7.48 (1H, d), 7.66 (1H, s), 7.86 (1H, d), 11.76 (1H, br).

Preparation Example 9: Synthesis of 3-(4-amino-3,5-dichlorobenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione (compound 9)

From 1.06 g (4.40 mmol) of 4-amino-3,5-dichlorobenzenesulfonamide and 1.34 g (4.40 mmol) of methyl 4-chloro-2-N-phenoxy-carbonylanthranilate in a similar manner to Preparation Example 1, 905 mg (yield 44%) of methyl 2-{[(4-amino-3,5-dichlorobenzenesulfonylamino)carbonyl]amino}-4-chlorobenzoate (property: colorless amorphous, PMR (δ ppm, DMSO- d_6): 3.87 (3H, s), 6.59 (2H, br), 7.22 (1H, dd), 7.72 (2H, s), 7.93 (1H, d), 8.24 (1H, d), 10.17 (1H, s)) was obtained.

Then, from the 905 mg (2.0 mmol) of the sulfonylurea product obtained, 660 mg (yield 82%: 2 steps) of the title compound was obtained in a similar manner.

Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 6.80 (2H, s), 7.12 (1H, s), 7.24 (1H, d), 7.86 (1H, d), 7.92 (2H, s), 11.63 (1H, br).

Preparation Example 10: Synthesis of 3-(3-amino-4-methylbenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione (compound 10)

From 960 mg (3.00 mmol) of 3-benzyloxycarbonylamino-4-methylbenzenesulfonamide and 1.14 g (3.00 mmol) of 4-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in

a similar manner to Preparation Example 8, 1.14 g (yield 62%) of 2-[[(3-benzyloxycarbonylamino-4-methylbenzenesulfonylamino)carbonyl]amino}-4-chlorobenzoic acid benzylester (property: colorless amorphous, PMR (δ ppm, DMSO- d_6): 2.30 (3H, s), 5.17 (2H, s), 5.36 (2H, s), 7.20 (1H, dd), 7.33-7.48 (11H, m), 7.63 (1H, d), 7.97 (1H, d), 8.11 (1H, s), 8.25 (1H, s), 9.27 (1H, s), 10.30 (1H, s), 12.20 (1H, br)) was obtained.

Then, from the 1.14 g (1.87 mmol) of the sulfonylurea product obtained, 190 mg (yield 27%: 2 steps) of the title compound was obtained in a similar manner. Property: colorless crystals, melting point: $>200^\circ\text{C}$ (decomposition), PMR (δ ppm, DMSO- d_6): 2.12 (3H, s), 5.47 (2H, s), 7.12 (1H, s), 7.16-7.25 (3H, m), 7.38 (1H, s), 7.85 (1H, d), 11.58 (1H, s).

Preparation Example 11: Synthesis of 3-[(3-carboxymethylaminophenyl)sulfonyl]-7-chloro-2,4(1H,3H)-quinazolinedione (compound 11)

From 1.62 g (5.65 mmol) of 3-t-butoxycarbonylmethylaminobenzenesulfonamide and 1.73 g (5.65 mmol) of methyl 4-chloro-2-N-phenoxy-carbonylanthranilate in a similar manner to Preparation Example 7, 209 mg (yield 9%: 4 steps) of the title compound was obtained. Property: colorless crystals, melting point: $>200^\circ\text{C}$ (decomposition), PMR (δ ppm, DMSO- d_6): 3.86 (2H, s), 6.88 (1H, s), 7.12 (1H, s), 7.24 (1H, d), 7.30-7.38 (3H, m), 7.86 (1H, d), 11.61 (1H, br).

Preparation Example 12: Synthesis of 3-(3-aminobenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione (compound 12)

From 3.5 g (12.9 mmol) of 3-t-butoxycarbonylaminobenzenesulfonamide and 3.9 g (12.8 mmol) of methyl 4-chloro-2-N-phenoxy-carbonylanthranilate in a similar manner to Preparation Example 7, 2.2 g (yield 49%: 4 steps) of the title compound was obtained.

Property: colorless crystals, melting point: $>200^{\circ}\text{C}$
(decomposition), PMR (δppm , $\text{DMSO}-d_6$): 5.72 (2H, s), 6.87
(1H, d), 7.12 (1H, s), 7.23-7.27 (2H, m), 7.33 (1H, s),
7.86 (1H, d), 11.61 (1H, s).

5 Preparation Example 13: Synthesis of 2-{3-[(7-chloro-
2,4(1H,3H)-quinazolinedione-3-
yl)sulfonyl]phenylaminocarbonyl}propionic acid (compound
13)

One hundred mg (0.28 mmol) of compound 12 was
10 dissolved in 5 ml of THF, to which 100 mg (1.0 mmol) of
succinic anhydride was added and heated to reflux for 3
hours. The reaction mixture was concentrated under
reduced pressure, and the crude product obtained was
crystallized from ethyl acetate-diethylether to obtain
15 120 mg (yield 96%) of the title compound. Property:
colorless crystals, melting point: $187-188^{\circ}\text{C}$, PMR (δppm ,
 $\text{DMSO}-d_6$): 2.54 (2H, d), 2.59 (2H, d), 7.12 (1H, s), 7.24
(1H, d), 7.59 (1H, t), 7.80 (1H, d), 7.86 (1H, d), 7.96
(1H, d), 8.41 (1H, s), 10.40 (1H, s), 11.63 (1H, br),
20 12.10 (1H, br).

Preparation Example 14: Synthesis of 3-{3-[(7-chloro-
2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]phenyl}acrylic
acid (compound 14)

From 1.54 g (5.44 mmol) of 3-(3-
25 aminosulfonyl)phenylacrylic acid t-butylester and 1.66 g
(5.44 mmol) of methyl 4-chloro-2-N-
phenoxy-carbonylanthranilate in a similar manner to
Preparation Example 7, 2.18 g (yield 81%) of methyl 2-
({[3-{3-t-butoxy-3-oxo-1-
30 propenyl)benzenesulfonylamino]carbonyl}amino)-4-
chlorobenzoate (property: colorless amorphous, PMR (δppm ,
 CDCl_3): 1.53 (9H, s), 3.95 (3H, s), 6.46 (1H, d), 7.05
(1H, d), 7.55 (1H, m), 7.57 (1H, d), 7.72 (1H, m), 7.93
(1H, m), 8.04 (1H, m), 8.27 (1H, s), 8.46 (1H, d), 11.05
35 (1H, br)) was obtained.

Then, from the 2.18 g (4.4 mmol) of the sulfonylurea

product obtained, 698 mg (yield 37%: 3 steps) of the title compound was obtained in a similar manner.

Property: colorless crystals, melting point: >200°C

(decomposition), PMR (δ ppm, DMSO- d_6): 6.65 (1H, d), 7.12 (1H, s), 7.25 (1H, d), 7.69 (1H, d), 7.72 (1H, t), 7.87 (1H, d), 8.12 (2H, q), 8.37 (1H, s), 11.64 (1H, s).

Preparation Example 15: Synthesis of 4-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]salicylic acid (compound 15)

From 1.0 g (3.66 mmol) of 4-t-butoxycarbonyl-3-hydroxybenzenesulfonamide and 1.12 g (3.66 mmol) of methyl 4-chloro-2-N-phenoxy-carbonylanthranilate in a similar manner to Preparation Example 7, 1.79 g (yield 100%) of methyl 2-{[(4-t-butoxycarbonyl-3-hydroxybenzenesulfonylamino)carbonyl]amino}-4-chlorobenzoate (property: colorless amorphous, PMR (δ ppm, DMSO- d_6): 1.57 (9H, s), 3.87 (3H, s), 7.14 (1H, d), 7.40-7.45 (2H, m), 7.85 (1H, d), 7.92 (1H, d), 8.32 (1H, d), 10.13 (1H, s), 10.82 (1H, s)) was obtained.

Then, from the 1.78 g (3.66 mmol) of the sulfonylurea product obtained, 370 mg (yield 25%: 3 steps) of the title compound was obtained in a similar manner. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 7.13 (1H, s), 7.26 (1H, d), 7.69 (1H, d), 7.87 (1H, d), 8.01 (1H, d), 11.67 (1H, s).

Preparation Example 16: Synthesis of 4-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]salicylic acid monosodium salt (compound 16)

Fifty mg (0.13 mmol) of compound 15 was suspended in about 1 ml of THF, to which 126 μ l of 1N sodium hydroxide in water was added dropwise. After confirming that the solution became homogeneous, 30 ml of water was added and lyophilized to obtain 52 mg of the amorphous title compound on a quantitative basis. Property: colorless amorphous, PMR (δ ppm, CD₃OD): 7.11 (1H, s), 7.19 (1H, d),

7.58 (1H, d), 7.63 (1H, s), 7.92 (1H, d), 8.03 (1H, d).

Preparation Example 17: Synthesis of 4-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]anthranilic acid (compound 17)

5 From 2.84 g (6.99 mmol) of 3-benzyloxycarbonylamino-4-t-butoxycarbonylbenzenesulfonamide and 2.67 g (6.99 mmol) of 4-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example 8, 3.74 g (yield 77%) of 2-[(3-benzyloxycarbonylamino-4-t-butoxycarbonylbenzenesulfonylamino)carbonyl]amino}-4-chlorobenzoic acid benzylester (property: colorless amorphous, PMR (δ ppm, DMSO- d_6): 1.54 (9H, s), 5.19 (2H, s), 5.34 (2H, s), 7.05 (1H, m), 7.34-7.58 (10H, m), 7.60 (1H, d), 7.90 (1H, d), 7.98 (1H, d), 8.50 (1H, br), 8.62 (1H, s), 10.00 (1H, br), 10.41 (1H, s)) was obtained.

15 Then, from the 3.74 g (5.39 mmol) of the sulfonylurea product obtained, 690 mg (yield 30%: 2 steps) of 4-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]anthranilic acid t-butylester was obtained in a similar manner, which was subjected to a similar debutylation reaction to obtain 503 mg (yield 84%) of the title compound. Property: colorless crystals, melting point: $>200^\circ\text{C}$ (decomposition), PMR (δ ppm, DMSO- d_6): 7.14 (1H, s), 7.18 (1H, d), 7.25 (1H, d), 7.59 (1H, s), 7.87 (1H, d), 7.89 (1H, d), 11.62 (1H, s).

Preparation Example 18: Synthesis of 4-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]anthranilic acid monosodium salt (compound 18)

30 Fifty mg (0.13 mmol) of compound 17 was suspended in about 1 ml of THF, to which 126 μl of 1N sodium hydroxide in water was added dropwise. After confirming that the solution became homogeneous, 30 ml of water was added and lyophilized to obtain 52 mg of the amorphous title compound on a quantitative basis. Property: colorless amorphous, PMR (δ ppm, DMSO- d_6): 7.11-7.22 (3H, m), 7.37 (1H, s), 7.83 (1H, d), 7.91 (1H, d).

Preparation Example 19: Synthesis of 3-(4-hydroxybenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione (compound 19)

From 1.50 g (7.03 mmol) of 4-allyloxybenzenesulfonyl isocyanate and 1.2 g (7.03 mmol) of 4-chloroanthranilic acid in a similar manner to Preparation Example 5, 1.5 g (yield 53%) of 3-(4-allyloxybenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione was obtained. From 500 mg (1.27 mmol) among this, 405 mg of the title compound (yield 90%) was obtained in a similar manner. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 6.98 (2H, d), 7.11 (1H, s), 7.23 (1H, d), 7.85 (1H, d), 8.00 (2H, d), 11.25 (1H, br).

Preparation Example 20: Synthesis of 4-[(2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]salicylic acid (compound 20)

From 618 mg (2.26 mmol) of 4-t-butoxycarbonyl-3-hydroxybenzenesulfonamide and 613 mg (2.26 mmol) of 2-N-phenoxycarbonylanthranilic acid methylester in a similar manner to Preparation Example 17, 792 mg (yield 78%) of methyl 2-[(4-t-butoxycarbonyl-3-hydroxybenzenesulfonylamino)carbonylamino]benzoate (property: colorless amorphous, PMR (δ ppm, CDCl₃): 1.60 (9H, s), 3.97 (3H, s), 7.09 (1H, t), 7.49-7.52 (2H, m), 7.65 (1H, d), 7.90 (1H, d), 8.01 (1H, dd), 8.33 (1H, d), 10.98 (1H, s), 11.18 (1H, s)) was obtained.

Then, from the 790 mg (1.75 mmol) of the sulfonylurea product obtained, 100 mg (yield 8%: 3 steps) of the title compound was obtained in a similar manner. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 7.13 (1H, d), 7.22 (1H, t), 7.63-7.69 (3H, m), 7.87 (1H, d), 8.01 (1H, d), 11.57 (1H, s).

Preparation Example 21: Synthesis of 5-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]salicylic acid (compound 21)

From 320 mg (1.17 mmol) of 3-t-butoxycarbonyl-4-hydroxybenzenesulfonamide and 447 mg (1.17 mmol) of 4-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example 17, 611 mg (yield 93%) of 2-{[(3-t-butoxycarbonyl-4-hydroxybenzenesulfonylamino)carbonylamino]-4-chlorobenzoic acid benzylester (property: colorless amorphous, PMR (δ ppm, CDCl_3): 1.62 (9H, s), 5.35 (2H, s), 7.01-7.05 (2H, m), 7.37-7.41 (5H, m), 7.96 (1H, d), 8.10 (1H, dd), 8.46-8.48 (2H, m), 10.99 (1H, s), 11.66 (1H, s)) was obtained.

Then, from the 611 mg (1.09 mmol) of the sulfonylurea product obtained, 114 mg (yield 33%: 3 steps) of the title compound was obtained in a similar manner. Property: colorless crystals, melting point: $>200^\circ\text{C}$ (decomposition), PMR (δ ppm, $\text{DMSO}-d_6$): 7.11 (1H, s), 7.19 (1H, d), 7.24 (1H, d), 7.86 (1H, d), 8.20 (1H, d), 8.56 (1H, s), 11.57 (1H, s).

Preparation Example 22: Synthesis of 3-(3-acetamide-4-methoxybenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione (compound 22)

From 500 mg (2.19 mmol) of 3-acetamide-4-methoxybenzenesulfonamide and 836 mg (2.19 mmol) of 4-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner Preparation Example 8, 812 mg (yield 70%) of 2-{[(3-acetylamino-4-methoxybenzenesulfonylamino)carbonylamino]-4-chlorobenzoic acid benzylester (property: colorless amorphous, PMR (δ ppm, $\text{DMSO}-d_6$): 2.12 (3H, s), 3.93 (3H, s), 5.36 (2H, s), 7.20 (1H, d), 7.24 (1H, d), 7.36-7.48 (5H, m), 7.69 (1H, d), 7.96 (1H, d), 8.24 (1H, d), 8.67 (1H, s), 9.39 (1H, s), 10.25 (1H, s), 12.11 (1H, br)) was obtained.

Then, from the 611 mg (1.09 mmol) of the sulfonylurea product obtained, 250 mg (yield 39%: 2 steps) of the title compound was obtained in a similar manner. Property: colorless crystals, melting point:

5 >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 2.12 (3H, s), 3.95 (3H, s), 7.12 (1H, s), 7.23 (1H, d), 7.30 (1H, d), 7.85 (1H, d), 7.89 (1H, d), 8.80 (1H, s), 9.42 (1H, s), 11.59 (1H, br)).

10 Preparation Example 23: Synthesis of 3-(3-amino-4-methoxybenzenesulfonyl)-7-chloro-2,4(1H,3H)-quinazolinedione (compound 23)

From 400 mg (1.40 mmol) of 3-t-butoxycarbonylamino-4-methoxybenzenesulfonamide and 533 mg (1.40 mmol) of 4-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example 17, 86 mg (yield 16%: 4 steps) of the title compound was obtained.

15 Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 3.81 (3H, s), 7.26-7.37 (5H, m), 7.77 (1H, s), 7.90 (1H, d), 7.94 (1H, d), 11.73 (1H, s)

20 Preparation Example 24: Synthesis of 7-chloro-3-(4-methoxy-3-methylsulfonylamino)benzenesulfonyl)-2,4(1H,3H)-quinazolinedione (compound 24)

From 500 mg (1.89 mmol) of 4-methoxy-3-methylsulfonylamino)benzenesulfonamide and 722 mg (1.89 mmol) of 4-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example 8, 888 mg (yield 83%) of 2-([(4-methoxy-3-methylsulfonylamino)benzenesulfonylamino]carbonyl)amino)-4-chlorobenzoic acid benzylester (property: colorless amorphous, PMR (δ ppm, DMSO- d_6): 2.12 (3H, s), 3.93 (3H, s), 5.36 (2H, s), 7.20 (1H, d), 7.24 (1H, d), 7.36-7.48 (5H, m), 7.69 (1H, d), 7.96 (1H, d), 8.24 (1H, s), 8.67 (1H, s), 9.39 (1H, s), 10.25 (1H, s), 12.11 (1H, br)) was obtained.

35 Then, from the 880 mg (1.55 mmol) of the

sulfonylurea product obtained, 620 mg (yield 85%: 2 steps) of the title compound was obtained in a similar manner. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 3.04 (3H, s), 3.94 (3H, s), 7.11 (1H, s), 7.23 (1H, d), 7.34 (1H, d), 7.86 (1H, d), 7.99 (1H, d), 8.10 (1H, s)).

Preparation Example 25: Synthesis of 4-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]-1-hydroxy-2-naphthylic acid (compound 25)

From 323 mg (1.00 mmol) of 3-t-butoxycarbonyl-4-hydroxy-1-naphthalenesulfonamide and 381 mg (1.00 mmol) of 4-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example 17, 447 mg (yield 73%) of 4-([(2-benzyloxycarbonyl-5-chloroanilino)carbonyl]amino)sulfonyl)-1-hydroxy-2-naphthalene carboxylic acid t-butylester (property: colorless amorphous, PMR (δ ppm, DMSO- d_6): 1.66 (9H, s), 5.34 (3H, s), 6.98 (1H, d), 7.35-7.48 (5H, m), 7.66 (1H, m), 7.81 (1H, m), 7.89 (1H, d), 8.37 (2H, m), 8.44 (1H, s), 8.71 (1H, d), 10.02 (1H, br), 12.52 (1H, br)) was obtained.

Then, from the 445 mg (0.72 mmol) of the sulfonylurea product obtained, 56 mg (yield 18%: 3 steps) of the title compound was obtained in a similar manner. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 7.08 (1H, s), 7.20 (1H, d), 7.63 (1H, t), 7.77 (1H, t), 7.84 (1H, d), 8.42 (1H, d), 8.51 (1H, d), 8.75 (1H, s), 11.57 (1H, s).

Preparation Example 26: Synthesis of 5-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]anthranilic acid (compound 26)

From 834 mg (2.05 mmol) of 4-benzyloxycarbonylamino-3-t-butoxycarbonylbenzenesulfonamide and 783 mg (2.05 mmol) of 4-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example 17, 1.18 g (yield 83%) of 2-([(4-benzyloxycarbonylamino-

3-t-butoxycarbonylbenzenesulfonylamino)carbonyl]amino}-4-chlorobenzoic acid benzylester (property: colorless amorphous, PMR (δ ppm, CDCl_3): 1.56 (9H, s), 5.22 (2H, s), 5.37 (2H, s), 7.04 (1H, dd), 7.33-7.42 (10H, m), 7.97 (1H, d), 8.14 (1H, d), 8.45 (1H, d), 8.60 (1H, d), 8.65 (1H, d), 11.01 (1H, s), 11.11 (1H, s)) was obtained.

Then, from the 1.17 g (1.69 mmol) of the sulfonylurea product obtained, 404 mg (yield 60%: 3 steps) of the title compound was obtained in a similar manner. Property: colorless crystals, melting point: $>200^\circ\text{C}$ (decomposition), PMR (δ ppm, $\text{DMSO}-d_6$): 6.89 (1H, d), 7.11 (1H, s), 7.23 (1H, d), 7.85 (1H, d), 7.98 (1H, d), 8.51 (1H, s), 11.51 (1H, s).

Preparation Example 27: Synthesis of 4-[(7-methoxy-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]anthranilic acid (compound 27)

From 500 mg (1.23 mmol) of 3-benzyloxycarbonylamino-4-t-butoxycarbonylbenzenesulfonamide and 460 mg (1.22 mmol) of 4-methoxy-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example 17, 15 mg (yield 3.1%: 4 steps) of the title compound was obtained. Property: colorless crystals, melting point: $>200^\circ\text{C}$ (decomposition), PMR (δ ppm, $\text{DMSO}-d_6$): 3.82 (3H, s), 6.58 (1H, s), 6.80 (1H, d), 7.16 (1H, d), 7.56 (1H, s), 7.80 (1H, d), 7.90 (1H, d), 11.49 (1H, s).

Preparation Example 28: Synthesis of (\pm)-7-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]-2-oxo-1H,3H-quinoline-3-carboxylic acid (compound 28)

From 400 mg (1.23 mmol) of (\pm)-3-t-butoxycarbonyl-2-oxo-1H,3H-quinoline-7-sulfonamide and 468 mg (1.23 mmol) of 4-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example 17, 649 mg (yield 86%) of 8-([(2-benzyloxycarbonyl-5-chloroanilino)carbonyl]amino)sulfonyl)-2-oxo-1,2,3,4-tetrahydro-3-quinolinecarboxylic acid t-butylester (property: colorless amorphous, PMR (δ ppm, CDCl_3): 1.32

(9H, s), 3.18-3.30 (2H, m), 3.54 (1H, m), 5.35 (2H, s), 6.85 (1H, m), 7.00 (1H, m), 7.35-7.39 (5H, m), 7.87-7.96 (3H, m), 8.47 (1H, m), 8.78 (1H, br), 10.92 (1H, br)) was obtained.

5 Then, from the 640 mg (1.04 mmol) of the sulfonylurea product obtained, 258 mg (yield 55%: 3 steps) of the title compound was obtained in a similar manner. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 3.23-3.31
10 (2H, m), 3.59 (1H, t), 7.07 (1H, d), 7.12 (1H, s), 7.25 (1H, d), 7.86 (1H, d), 7.96 (1H, d), 7.98 (1H, d), 10.84 (1H, s), 11.60 (1H, s).

Preparation Example 29: Synthesis of (\pm)-6-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]-3-oxo-1,4-benzoxadine-2-carboxylic acid (compound 29)
15

From 300 mg (0.91 mmol) of (\pm)-2-t-butoxycarbonyl-3-oxo-1,4-benzoxadine-6-sulfonamide and 349 mg (0.91 mmol) of 4-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example
20 17, 417 mg (yield 74%) of 5-({[(2-benzyloxycarbonyl-5-chloroanilino)carbonyl]amino}sulfonyl)-3-oxo-3,4-dihydro-2H-1,4-benzoxadine-2-carboxylic acid t-butylester (property: colorless amorphous, PMR (δ ppm, DMSO- d_6): 1.29 (9H, s), 5.37 (2H, s), 5.42 (2H, s), 7.19-7.26 (2H, m),
25 7.37-7.57 (7H, m), 7.97 (1H, d), 8.25 (1H, d), 10.27 (1H, s), 11.25 (1H, s), 12.22 (1H, br)) was obtained.

Then, from the 417 mg (0.68 mmol) of the sulfonylurea product obtained, 100 mg (yield 32%: 3 steps) of the title compound was obtained in a similar
30 manner. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 5.47 (1H, s), 7.11 (1H, s), 7.24 (1H, d), 7.29 (1H, d), 7.76 (1H, s), 7.78 (1H, d), 7.86 (1H, d), 11.25 (1H, s), 11.62 (1H, s).

Preparation Example 30: Synthesis of 4-[(7-hydroxy-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]anthranilic acid (compound 30)

From 620 mg (1.53 mmol) of 3-benzyloxycarbonylamino-4-t-butoxycarbonylbenzenesulfonamide and 550 mg (1.51 mmol) of 4-hydroxy-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example 17, 25 mg (yield 4%: 4 steps) of the title compound was obtained. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 6.48 (1H, s), 6.61 (1H, d), 7.14 (1H, d), 7.51 (1H, s), 7.70 (1H, d), 7.90 (1H, d), 10.80 (1H, s), 11.39 (1H, s).

Preparation Example 31: Synthesis of 4-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]-2-N-propionylantronic acid (compound 31):

Eight hundred and forty mg (1.86 mmol) of compound 17 was dissolved in 8 ml of 1,4-dioxane, to which 240 μ l (2.79 mmol) of propionyl chloride was added dropwise and stirred overnight at 60°C. Excess water was added to the reaction mixture and then extracted with ethyl acetate. The organic layer was washed, dried, and concentrated to obtain the crude product of 4-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]-2-N-propionylantronic acid t-butylester. After the crude product obtained was stirred in 3 ml of trifluoroacetic acid at room temperature for 1 hour, the reaction mixture was concentrated under reduced pressure to obtain a crude product, which was washed with diethylether to obtain 400 mg (yield 48%: 2 steps) of the title compound. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO- d_6): 1.10 (3H, t), 2.45 (2H, dd), 7.11 (1H, s), 7.24 (1H, d), 7.85 (1H, d), 7.88 (1H, d), 8.17 (1H, d), 9.18 (1H, s), 11.07 (1H, s), 11.63 (1H, s).

Preparation Example 32: Synthesis of 4-[(6-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]anthranilic acid (compound 32)

From 300 mg (0.74 mmol) of 3-benzyloxycarbonylamino-4-t-butoxycarbonylbenzenesulfonamide and 310 mg (0.81 mmol) of 5-chloro-2-N-phenoxy-carbonylanthranilic acid benzylester in a similar manner to Preparation Example 17, 75 mg (yield 26%: 4 steps) of the title compound was obtained. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO-d₆): 7.13-7.20 (2H, m), 7.56 (1H, s), 7.72 (1H, d), 7.82 (1H, s), 7.90 (1H, d), 11.68 (1H, s).

Preparation Example 33: Synthesis of 4-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]-2-N-methanesulfonylanthranilic acid (compound 33)

From 200 mg (0.44 mmol) of compound 17 in a similar manner to Preparation Example 3, 81 mg of 4-[(7-chloro-2,4(1H,3H)-quinazolinedione-3-yl)sulfonyl]-2-N-methanesulfonylanthranilic acid t-butylester was obtained, which was similarly subjected to a debutylation reaction to obtain 53 mg (yield 25%: 2 steps) of the title compound. Property: colorless crystals, melting point: >200°C (decomposition), PMR (δ ppm, DMSO-d₆): 3.24 (3H, s), 7.11 (1H, s), 7.25 (1H, d), 7.85-7.91 (2H, m), 8.23 (1H, d), 8.39 (1H, s), 11.05 (1H, br), 11.70 (1H, s).

Preparation Example 34: Synthesis of 3-(3-aminobenzenesulfonyl)-7-chloro-2,4(1H,3H)quinazolinedione methanesulfonic acid salt (compound 34)

2.15 g (6.10 mmol) of compound 12 was dissolved in 65 ml of THF, to which 0.4 ml of methanesulfonic acid was added. To this solution was added 200 ml of ether, and a precipitate that deposited was filtered to obtain 2.59 g (yield 95%) of the title compound. Property: colorless amorphous, PMR (δ ppm, DMSO-d₆): 2.35 (3H, s), 6.98 (1H, d), 7.12 (1H, m), 7.25 (1H, m), 7.34 (2H, s), 7.43 (1H,

m), 7.86 (1H, s), 11.64 (1H, s).

Example 1: Evaluation of inhibitory activity of test compounds to human chymase

Human heart chymase was purified according to the
5 method of Urata et al. (J. Biol. Chem. 265: 22348, 1990).
The inhibitory activity of the compound of the present
invention was determined as follows. Purified enzyme was
diluted with 0.1 M Tris-HCl buffer (pH 7.5) containing 1
10 M sodium chloride and 0.01% Triton X-100 to appropriate
concentrations. Suc-Ala-Ala-Pro-Phe-MCA (Peptide
Institute Inc.) was dissolved in 10 mM dimethyl sulfoxide
(hereinafter referred to as DMSO) and diluted 20-fold
with 0.1 M Tris-HCl buffer (pH 7.5) containing 1 M sodium
15 chloride and 0.01% Triton X-100 to an appropriate
concentration to prepare substrate solution.

5 µl of the test sample in DMSO was added to 75 µl
of the enzyme solution and preincubated at 30°C for 10
minutes. Then, 20 µl of the substrate solution was added
to the test sample-enzyme mixture, and incubated at 30°C.
20 Ten minutes later, 50 µl of 30% acetic acid was added to
stop the enzymatic reaction, and the amount of AMC formed
was determined using a fluorophotometer. At the same
time, 5 µl of DMSO in stead of the test sample was added
and reacted simultaneously as a control. The inhibitory
25 activity to human chymase was calculated based on the
value of the control, and then the inhibition percentage
and the 50% inhibition concentration (IC₅₀) were
determined.

The IC₅₀ values of representative compounds are
30 shown in Table 1.

Table 1

Preparation Example No.	IC ₅₀ value (μ M)	t _{1/2} (min)
1	0.36	78
2	0.14	175
8	0.035	29
10	0.17	167
12	0.44	249
13	0.3	97
16	0.84	>240
17	0.14	260
18	0.14	103
21	0.34	-
22	0.3	104
24	0.32	79
27	4.0	263
29	1.7	>240
32	1.5	74
34	0.36	709

Example 2: Stability in human plasma

2 μ l of 1 mM test sample in DMSO was added to 198 μ l
 5 of 50% of human plasma solution in a 50 mM sodium
 phosphate buffer (pH 7.2) and incubated at 30°C. At 0,
 5, and 15 minutes, 800 μ l of acetonitrile was added to
 the test sample-plasma mixture, which was mixed and
 deproteinized. The supernatant obtained by
 10 centrifugation (12,000 rpm, one minute) was diluted with
 the same volume of distilled water, and the intact
 compound in the solution was determined by HPLC analysis.

The half life (t_{1/2}) of the test sample in the plasma
 solution was calculated from the recovery at each time
 15 point using an exponential regression analysis. The
 plasma half life (t_{1/2}) for representative compounds is
 shown in Table 1.

Example 3: Aorta lipid deposition model in hamster

An aorta lipid deposition model was induced by a
 20 high-cholesterol diet. A high-cholesterol diet was
 prepared by adding 0.5% cholesterol and 10% coconut oil
 (KBT Oriental Co.) to a standard rodent feed containing
 5.1% fat and 0.07% cholesterol (KBT Oriental Co.). 8-
 week-old male Golden Syrian hamsters (KBT Oriental Co.)

(100-130 g) were given the high-cholesterol diet for 12 weeks to induce lipid deposition in the aorta. A group to which a standard rodent feed was given for 12 weeks was used as the control.

5 On week 12 after the start of the high-cholesterol diet, total cholesterol, low density lipoprotein (LDL) cholesterol, and high density lipoprotein (HDL) cholesterol levels in the peripheral blood and chymase-like activity in the aorta was determined. Chymase-like
10 activity was measured using Ang I as the substrate, and expressed by subtracting the activity inhibited by aprotinin from the activity inhibited by chymostatin (M. Akasu, et al., Hypertension 32: 514-20, 1998, M. Ihara, et al., Hypertension 33: 1399-405, 1999).

15 Lipid deposition in the aorta was evaluated by harvesting the aorta on week 12 after the start of the high-cholesterol diet and performing a histopathological analysis. Namely, the ascending aorta from its junction with the heart to the middle part was removed, washed in
20 an ice-cold saline, and then 3-5 mm segments containing the aortic cusp region were cryopreserved in Tissue-Tek O.C.T. Compound (Miles Inc.). Then, the frozen sections of 6 μ m were prepared, fixed in 10% formalin for 10 minutes, washed with distilled water, and stained with
25 the Oil red O welding (Muto Pure Chemicals) at 60°C for 5 minutes.

 Then, the sections were washed with 60% isopropanol and distilled water, and counterstained with hematoxylin for 2 minutes. After washing with 1/4 saturated LiCO₃,
30 lipid deposition was evaluated by microscopic observation. Also, the area of the lipid deposition region (the region stained to an orange color with Oil red O) was quantitated on the histological pictures by NIH Image software ver. 1.61.

35 Result:

 In the aorta of high cholesterol diet-treated hamster, a conspicuous lipid deposition was observed in

the intimal region. In this model (n=6) as compared to the control group (n=6), a marked increase in the lipid deposition area in the aortic cusp region was observed (Figure 1).

5 Also, there was a significant increase in plasma levels of total cholesterol, LDL cholesterol and HDL cholesterol in this model as compared to the control group (their respective values were 4.16 ± 0.36 vs. 12.59 ± 1.01 mmol/L, 1.15 ± 0.26 vs. 5.48 ± 0.67 mmol/L and
10 1.86 ± 0.08 vs. 3.62 ± 0.10 mmol/L, n=6 each, $p < 0.01$).

 Furthermore, there was a significant increase in the chymase-like activity in the aorta in this model as compared to the control group (19.2 ± 2.6 vs. 9.3 ± 1.2 nmol/min/g wet tissue, respectively, n=6 each, $p < 0.01$).

15 A positive correlation was observed between the chymase-like activity and total cholesterol or LDL cholesterol levels (Figure 2).

 Taken together, these results demonstrate that chymase is involved in lipid deposition induced by a high
20 cholesterol exposure.

Example 4: Construction of transgenic mice that express human chymase at a high level

 Transgenic (Tg) mice that express human chymase at a high level were constructed according to the method as
25 described (Zokuseikagakujikkenkoza (Sequel to Biochemistry Experimental Series) 1, Idenshikenkyuhou (Gene Study Method) III, edited by the Japanese Biochemical Society). Briefly, a transgene was generated in which cDNA (J. Biol. Chem. 266: 17173, 1991) encoding
30 human chymase was placed under the control of chicken β actin promoter and the cytomegalovirus immediate early gene promoter. On the next day of mating, fertilized eggs were collected from the oviduct of female mice, and then the above transgene solution was injected into the
35 male pronucleus of the fertilized egg using a thin glass pipet. Fifteen to thirty of these fertilized eggs were transplanted into the oviduct of pseudopregnant female

mice, and about 20 days later, the eggs were allowed to undergo the natural or cesarean birth. The newborn mice were bred, and at about 4-week old, DNA was extracted from part of the tail portion, and the presence of the DNA of the introduced gene was searched using Southern blot method (Current Protocols in Molecular Biology, Wiley). The expression of human chymase in each tissue of Tg mice was investigated by Northern blot and Western blot analyses.

Result:

Tg mice obtained were subjected to spontaneous mating to obtain the offspring (F1) mice, in which the homozygous Tg mice that express human chymase were lethal. On the other hand, naturally born heterozygous Tg mice were found to express human chymase by Northern blot and Western blot in the heart, the blood vessel, the skin, the liver, the lung, and the brain (n=3). The body weight of these Tg mice were slightly smaller than the control mice (wild type littermates) (87% for males and 80% for females, each n=6), and they had hypotrichosis and Leukocytosis (Tg mice had $13300 \pm 3600 \mu\text{l}$ [n=14] relative $7700 \pm 2200 \mu\text{l}$ for the control mice [n=15], $p < 0.001$, t-test). On the other hand, blood pressure of 12-week old Tg mice was about the same level as the control mice ($116 \pm 15 \text{ mmHg}$ [n=10] and $108 \pm 9 \text{ mmHg}$ [n=10] in Tg mice and the control mice, respectively).

Example 5: Effect of the high-cholesterol diet on human chymase Tg mice

Effect of the high-cholesterol diet on the human chymase Tg mice (8-week old each) described in Example 4 was investigated. Preparation of the high-cholesterol diet, the evaluation of lipid deposition in the aorta, and the determination of chymase activity in the aorta were performed in the methods described in Example 6 below.

Result:

Chymase activity in the aorta of the human chymase Tg mice was significantly higher than that of the control mice. (Figure 3, n=6, p<0.05, t-test). Also, when a high-cholesterol diet is given, a significant increase in the lipid deposition area was observed in the human chymase Tg mice regardless of the sex (Figure 4, n=3).

The fact that high-cholesterol diet induced lipid deposition in the blood vessel in human chymase Tg mice but not in the control mice indicates the possibility that chymase inhibitor would be effective against lipid deposition on the blood vessel.

Example 6: Effect of chymase inhibitor in the aortic lipid deposition model in hamster

An aortic lipid deposition model was generated as described in Example 3 and used as the high-cholesterol diet group (n=6). A group that received the standard rodent diet for 12 weeks was used as the control group (n=6). The compound obtained in Preparation Example 18 (compound 18) was administered orally to a group that received high-cholesterol diet at a dose of 100 mg/kg/day every day for the same 12 weeks (n=6).

The effect on lipid deposition by the chymase inhibitor was evaluated by harvesting the aorta on week 12 after the start of the high-cholesterol diet and performing a histopathological analysis. Namely, the ascending aorta from its junction with the heart to the middle part was removed, washed in an ice-cold saline, and then using 3-5 mm segments containing the aortic cusp region were cryopreserved in Tissue-Tek O.C.T. Compound (Miles Inc.). Then, the frozen sections of 6 μ m were prepared, fixed in 10% formalin for 10 minutes, washed with distilled water, and then stained with the Oil red O welding (Muto Pure Chemicals) at 60°C for 5 minutes.

Then, the sections were washed with 60% isopropanol and distilled water, and counterstained with hematoxylin for 2 minutes. After washing with 1/4 saturated LiCO₃, lipid deposition was evaluated by microscopic

observation. Also, the area of the lipid deposition region (the region stained to an orange color with Oil red O) was quantitated on the histological pictures by NIH Image software ver. 1.61.

5 Result:

In the microscopic examination of the aortic cusp 12 weeks after the start of the high-cholesterol diet, a conspicuous lipid deposition was observed in the intimal region of the aorta of the hamsters that received the high-cholesterol diet, but lipid deposition had completely disappeared in the aorta of the group that received compound 18. Furthermore, the result in which the area of the lipid deposition region was determined is shown in Figure 1. The hamsters that received the high-cholesterol diet showed a marked increase in the lipid deposition area in the aortic cusp region as compared to the control group, and the oral administration of compound 18 significantly suppressed the increase in the lipid deposition area.

The finding that compound 18 improves lipid deposition in the high cholesterol-diet model indicates that chymase inhibitor ameliorates the abnormal vascular function to normal, and that chymase inhibitor is useful for the treatment of new diseases accompanied by abnormal vascular function in which lipid deposition in the blood vessel is involved.

Formulation Example 1: Preparation of tablets

One hundred grams of compound 1 was mixed with 22.5 g of microcrystalline cellulose and 2.5 g of magnesium stearate, which was pressed into tablets by a single punch press to formulate tablets of 9 mm in diameter and 250 mg in weight that contained 200 mg per tablet of compound 1.

Formulation Example 2: Preparation of granules

Thirty grams of compound 1 was mixed well with 265 g of lactose and 5 g of magnesium stearate, which was compression molded, communicated, sized, and filtered to

prepare satisfactory 10% granules of 20-50 mesh.

Formulation Example 3: Preparation of rectal
suppositories

5 Witepsol H-15 (manufactured by Dynamit Nobel) was
heat-melted, to which compound 1 was added to a
concentration of 12.5 mg/ml, mixed into homogeneity.
Then this was injected into the die for rectal
suppositories in 2 ml portions, and cooled to obtain
rectal suppositories containing 25 mg/tablet of compound
10 1.

INDUSTRIAL APPLICABILITY

15 According to the present invention, the effect of a
chymase inhibitor of suppressing lipid deposition in the
blood vessel can effectively prevent or treat diseases
accompanied by abnormal vascular function.

CLAIMS

1. A preventive or therapeutic agent for diseases accompanied by abnormal vascular function in which lipid deposition in the blood vessel is involved, said agent
5 comprising a chymase inhibitor as an active ingredient.


2. The preventive or therapeutic agent according to claim 1, wherein the diseases accompanied by abnormal vascular function in which lipid deposition in the blood vessel is involved are arteriosclerosis, cardiac acute
10 coronary syndrome, restenosis after percutaneous transluminal coronary angioplasty, obstructive arteriosclerosis, obstructive thrombotic vasculitis, atherosclerosis, cerebral infarction, intermittent claudication, lower limb gangrene, renal vascular
15 hypertension, renal arterial aneurysm, and renal infarction.

3. A preventive or therapeutic pharmaceutical composition for diseases accompanied by abnormal vascular function, wherein the chymase inhibitor is blended at an
20 amount that suppresses lipid deposition in the blood vessel.

4. The preventive or therapeutic pharmaceutical composition according to claim 3, wherein the diseases accompanied by abnormal vascular function are
25 arteriosclerosis, cardiac acute coronary syndrome, restenosis after percutaneous transluminal coronary angioplasty, obstructive arteriosclerosis, obstructive thrombotic vasculitis, atherosclerosis, cerebral infarction, intermittent claudication, lower limb
30 gangrene, renal vascular hypertension, renal arterial aneurysm, and renal infarction.

5. A suppressing agent of lipid deposition in the blood vessel comprising a chymase inhibitor as an active ingredient.

35 6. The preventive or therapeutic agent according to claim 1 or 2, wherein said chymase inhibitor is a quinazoline derivative represented by the formula (1):



5

wherein, the ring A represents an aryl ring,

10 R^1 represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower alkyl group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;

20 R^2 and R^3 , which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 4 carbons that may be substituted, a halogen atom, a hydroxy group, a lower alkoxy group having 1 to 4 carbons, an amino group, a lower alkylamino group having 1 to 4 carbons that may be substituted, a lower aralkylamino group having 7 to 10 carbons that may be substituted, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an

35

T08250-09269666


aromatic carboxylic acid that may be substituted with a
carboxylic group, an amino group acylated with a
heteroaromatic carboxylic acid that may be substituted
with a carboxylic group, an amino group sulfonylated with
5 a lower alkanesulfonic acid having 1 to 4 carbons that
may be substituted with a carboxylic group, an amino
group sulfonylated with an aromatic sulfonic acid that
may be substituted with a carboxylic group, an amino
group sulfonylated with a heteroaromatic sulfonic acid
10 that may be substituted with a carboxylic group, or a
carboxylic group; or

when the ring A is a benzene ring, R^1 and R^2 ,
together with the benzene ring to be substituted, may
form a fused heterocyclic ring that may be substituted
15 with a carboxylic acid, and a carbon atom in said fused
heterocyclic ring may form a carbonyl group wherein R^3 is
as defined above; and

X represents a hydrogen atom, a lower alkyl group
having 1 to 4 carbons, a lower alkoxy group having 1 to 4
20 carbons, a halogen atom, a hydroxy group, an amino group,
or a nitro group;

or a pharmaceutically acceptable salt thereof.

7. The pharmaceutical composition according to
claim 3 or 4, wherein said chymase inhibitor is a
25 quinazoline derivative represented by the formula (1):



wherein, the ring A represents an aryl ring,

R^1 represents a hydroxy group, an amino group, or a
lower alkylamino group having 1 to 4 carbons that may be
35 substituted with a carboxylic group, a lower aralkylamino
group having 7 to 10 carbons that may be substituted with
a carboxylic group, an amino group acylated with a lower

5
10
15
20
25
30
35


when the ring A is a benzene ring, R^1 and R^2 ,

together with the benzene ring to be substituted, may form a fused heterocyclic ring that may be substituted with a carboxylic acid, and a carbon atom in said fused heterocyclic ring may form a carbonyl group wherein R³ is as defined above; and

X represents a hydrogen atom, a lower alkyl group having 1-4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;

or a pharmaceutically acceptable salt thereof.

8. The suppressing agent of lipid deposition according to claim 5, wherein said chymase inhibitor is a quinazoline derivative represented by the formula (1):



wherein, the ring A represents an aryl ring,

R¹ represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower

alkyl group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;

5 R^2 and R^3 , which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 4 carbons that may be substituted, a halogen atom, a hydroxy group, a lower alkoxy group having 1 to 4 carbons, an amino group, a lower alkylamino group having 1 to 4 carbons that may be substituted, a lower
10 aralkylamino group having 7 to 10 carbons that may be substituted, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a
15 carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino
20 group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, or a carboxylic group; or


25 when the ring A is a benzene ring, R^1 and R^2 , together with the benzene ring to be substituted, may form a fused heterocyclic ring that may be substituted with a carboxylic acid, and a carbon atom in said fused heterocyclic ring may form a carbonyl group wherein R^3 is
30 as defined above; and

X represents a hydrogen atom, a lower alkyl group having 1 to 4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;
35 or a pharmaceutically acceptable salt thereof.

ABSTRACT

There is provided a preventive or therapeutic agent
for diseases accompanied by abnormal vascular function in
which lipid deposition in the blood vessel is involved,
said agent comprising a chymase inhibitor as an active
ingredient.

As the chymase inhibitor, a quinazoline derivative
represented by the following formula is used.



In the above formula, the ring A represents an
aromatic ring.

Fig.1





Fig.2

A



B




Fig. 3





Fig. 4



Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Japanese Language Declaration (日本語宣言書)

私は、米国法典第35編119条(a)-(d)項又は365条(b)項に基づき下記の、米国外の国の少なくとも一カ国を指定している特許協力条約365(a)項に基づく国際出願、又は外国での特許出願もしくは発明者証の出願についての外国優先権をここに主張するとともに、優先権を主張している、本出願の前に出願された特許または発明者証の外国出願を以下に、枠内をマークすることで、示しています。

Prior Foreign Application(s)

外国での先行出願

11-311257 (Pat. Appln.)

Japan

(Number)
(番号)

(Country)
(国名)

(Number)
(番号)

(Country)
(国名)

I hereby claim foreign priority under Title 35, United States Code, Section 119 (a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Priority Not Claimed
優先権主張なし

1/November/1999

(Day/Month/Year Filed)
(出願年月日)

(Day/Month/Year Filed)
(出願年月日)

私は、第35編米国法典119条(e)項に基づいて下記の米国外の特許出願規定に記載された権利をここに主張いたします。

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below.

(Application No.)
(出願番号)

(Filing Date)
(出願日)

(Application No.)
(出願番号)

(Filing Date)
(出願日)

私は、下記の米国法典第35編120条に基づいて下記の米国外の特許出願に記載された権利、又は米国外を指定している特許協力条約365条(c)に基づき権利をここに主張します。また、本出願の各請求範囲の内容が米国法典第35編112条第1項又は特許協力条約で規定された方法で先行する米国外の特許出願に開示されていない限り、その先行米国外出願提出日以降で本出願書の日本国内または特許協力条約国際提出日までの期間中に入手された、連邦規則法典第37編1条56項で定義された特許資格の有無に関する重要な情報について開示義務があることを認識しています。

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s), or 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code Section 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of application.

(Application No.)
(出願番号)

(Filing Date)
(出願日)

(Status: Patented, Pending, Abandoned)
(現況: 特許許可済、係属中、放棄済)

(Application No.)
(出願番号)

(Filing Date)
(出願日)

(Status: Patented, Pending, Abandoned)
(現況: 特許許可済、係属中、放棄済)

私は、私自身の知識に基づいて本宣言書中で私が行なう表明が真実であり、かつ私の入手した情報と私の信じているところに基づき表明が全て真実であると信じていること、さらに故意になされた虚偽の表明及びそれと同等の行為は米国法典第18編第1001条に基づき、罰金または拘禁、もしくはその両方により処罰されること、そしてそのような故意による虚偽の声明を行えば、出願した、又は既に許可された特許の有効性が失われることを認識し、よってここに上記のごとく宣誓を致します。

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Declaration and Power of Attorney For Patent Application

特許出願宣言書及び委任状

Japanese Language Declaration

日本語宣言書

下記の氏名の発明者として、私は以下の通り宣言します。

As a below named inventor, I hereby declare that:

私の住所、私書箱、国籍は下記の私の氏名の後に記載された通りです。

My residence, post office address and citizenship are as stated next to my name.

下記の名称の発明に関して請求範囲に記載され、特許出願している発明内容について、私が最初かつ唯一の発明者（下記の氏名が一つの場合）もしくは最初かつ共同発明者であると（下記の名称が複数の場合）信じています。

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

BLOOD VESSEL LIPID DEPOSITION-PREVENTIVE

AGENT COMPRISING CHYMASE-INHIBITOR

上記発明の明細書（下記の欄でx印がついていない場合は、本書に添付）は、

the specification of which is attached hereto unless the following box is checked:

☐ 月 日に提出され、米国出願番号または特許協定条約国際出願番号を _____ とし、
(該当する場合) _____ に訂正されました。

☐ was filed on November 1, 2000
as United States Application Number or
PCT International Application Number
PCT/JP00/07706 and was amended on
_____ (if applicable).

私は、特許請求範囲を含む上記訂正後の明細書を検討し、内容を理解していることをここに表明します。

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

私は、連邦規則法典第37編第1条56項に定義されたとおり、特許資格の有無について重要な情報を開示する義務があることを認めます。

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Japanese Language Declaration (日本語宣言書)

委任状: 私は下記の発明者として、本出願に関する一切の
手続きを米特許商標局に対して遂行する弁理士または代理人
として、下記の者を指名いたします。(弁理士、または代理
人の氏名及び登録番号を明記のこと)

POWER OF ATTORNEY: As a named inventor, I hereby appoint
the following attorney(s) and/or agent(s) to prosecute this
application and transact all business in the Patent and Trademark
Office connected therewith (list name and registration number)

William L. Mathis 17,337
Peter H. Smolka 15,913
Robert S. Swecker 19,885
Platon N. Mandros 22,124
Benton S. Duffett, Jr. 22,030
Joseph R. Magnone 24,239
Norman H. Stepno 22,716
Ronald L. Grudziecki 24,970
Frederick G. Michaud, Jr. 26,003
Alan E. Kopecki 25,813
Regis E. Slutter 26,999
Samuel C. Miller, III 27,360
Ralph L. Freeland, Jr. 16,110

Robert G. Mukai 28,531
George A. Hovanec, Jr. 28,223
James A. LaBarre 28,632
E. Joseph Gess 28,510
R. Danny Huntington 27,903
Eric H. Weisblatt 30,505
James W. Peterson 26,057
Teresa Stanek Rea 30,427
Robert E. Krebs 25,885
William C. Rowland 30,888
T. Gene Dillahunt 25,423
Patrick C. Keane 32,858
Bruce J. Boggs, Jr. 32,344

William H. Benz 25,952
Peter K. Skiff 31,917
Richard J. McGrath 29,195
Matthew L. Schneider 32,814
Michael G. Savage 32,596
Gerald F. Swiss 30,113
Michael J. Ure 33,089
Charles F. Wieland III 33,096
Bruce T. Wieder 33,815
Todd R. Walters 34,040

書類送付先

Send Correspondence to:

Ronald L. Grudziecki
BURNS, DOANE, SWECKER & MATHIS, L.L.P.
P.O. Box 1404
Alexandria, Virginia 22313-1404

直接電話連絡先: (名前及び電話番号)

Direct Telephone Calls to: (name and telephone number)

Ronald L. Grudziecki
at (703) 836-6620

唯一または第一発明者名	Full name of sole or first inventor	
	Harukazu Fukami	
発明者の署名	Inventor's signature	Date
	Harukazu Fukami	June 11, 2001
住所	Residence	
	Kyoto-shi, Kyoto, Japan	
国籍	Citizenship	
	Japanese	
私書箱	Post Office Address	
	36, Kishoin-Shimadezaike-cho, Minami-ku, Kyoto-shi, Kyoto 601-8373, Japan	
第二共同発明者	Full name of second joint inventor, if any	
	Hidenori Urata	
第二共同発明者	Second inventor's signature	Date
	Hidenori Urata	June 11, 2001
住所	Residence	
	Fukuoka-shi, Fukuoka, Japan	
国籍	Citizenship	
	Japanese	
私書箱	Post Office Address	
	18-3, Minamikatae 5-chome, Jonan-ku, Fukuoka-shi, Fukuoka 814-0143, Japan	

(第三以降の共同発明者についても同様に記載し、署名をす
ること)

(Supply similar information and signature for third and subsequent
joint inventors.)